City of Santa Fe



CITY CLERK'S OFFICE

Agendante 9/1/14 TIMF, 7:444

SERVEU BY Melissa Mel)onald

RECEIVED BY

Santa Fe River Commission Agenda
Thursday, September 8, 2016 (Round House Room), 6 pm to 8 pm
City Offices at the Market Station Building at the Rail yard
500 Market Street, Suite 200, Santa Fe, NM
505-955-6840

- 1. ROLL CALL
- 2. APPROVAL OF AGENDA
- 3. APPROVAL OF MINUTES FROM JULY 14, 2016 & AUGUST 11, 2016
- 4. Communication from other Agencies/Committees
- 5. INFORMATION/DISCUSSION/ACTION:
 - a) Project Update Review (Melissa McDonald)
 - b) River Parkway Benches and Bike Rack / PNM Power Up Grant Update (Raquel Thompson)
 - c) Friends of Santa Fe River Walk Update (Anna Hansen)
 - d) SF River Watershed Study Vegetation Mapping Report (Melissa McDonald)
 - e) SUB-COMMITTEE BREAKOUT SESSION -Working meeting
 - Outdoor Economy
 - Promoting a Living River
 - Watershed Revitalization
 - Species Resiliency
- 6. MATTERS FROM COMMISSIONERS
- 7. MATTERS FROM STAFF
- 8. CITIZENS' COMMUNICATION FROM THE FLOOR
- 9. ADJOURN

Next Scheduled for the River Commission is October 13, 2016
Captions & Packet Material are due by October 5, 2016
Persons with disabilities in need of accommodations, contact the City Clerk's office at (505) 955-6521 five (5) working days prior to the meeting date.

Santa Fe River Commission Meeting Index September 8, 2016

Title	Description	
Cover Sheet		0
Call to Order	Vice Chair Bove called the meeting of the Santa Fe River Commission to order at 6:01 p.m. at 500 Market Station (Railyard Station), Santa Fe, NM.	
Roll Call	A quorum was established will roll call.	1
Approval of the Agenda	Ms. Hansen moved to approve the agenda as presented with a second from Ms. Doremus which passed by voice vote.	1
Approval of Minutes from July 14, 2016 & August 11, 2016	Ms Doremus moved to approve the minutes of July 14, 2016 as presented with a second from Ms. Isaacson which passed by voice vote.	1
	Mr. Jacobi moved to approve the minutes of August 11, 2016 as presented with a second from Ms. Hansen which passed by voice vote.	1
Communication from other Agencies/Committees	Discussion Only	1
Information/Discussion/Action		1
a.)Project Update Review	·	2,3
b.) River Parkway Benches and Bike Rack/PNM Power Up Grant Update c.) Friends of Santa Fe River Walk Update	Disauration Only	3
d) SF River Watershed Study Vegetation Mapping Report e) Sub-Committee Break Out Session- Working	Discussion Only	3
Meeting Outdoor Economy Promoting a Living River Watershed Revitalization		3
Species Resiliency	The standard Resident Annual to show name position	3,4
Matters From Commissioners	Ms. Hansen moved to allow Ms. McDonald to share name, position on the committee, addresses and phone numbers and length of terms to send out with a second from Mr. Patorni which passed by voice vote.	J ₁ 4
Matters from Staff	Discussion Only	4
Citizen's Communication From the Floor	Discussion Only	4
Adjourn	Mr. Jacobi moved to adjourn the meeting at 7:22 p.m. with a second from Mr. Pierpont which passed by voice vote.	4
Signature Page		4

Santa Fe River Commission Meeting Minutes-September 8, 2016 500 Market Street Santa Fe, New Mexico 6:00 p.m.-8:00 p.m.

CALL TO ORDER

Vice Chair Bové called the meeting of the Santa Fe River Commission to order at 6:01 p.m. at 500 Market Station (Railyard Station), Santa Fe, NM. A quorum was established will roll call.

1. ROLL CALL

Present

Phil Bové, Vice Chair Luke Pierpont Anna Hansen F.M. Patomi Jerry Jacobi Zoe Isaacson Dale Doremus

Not Present/Excused

John R. Buchser, Chair Emile Sawyer

Others Present

Alan Hook, City of Santa Fe Water Division Andy Otto, Executive Director Santa Fe Watershed Association Raquel Baca-Tompson, Santa Fe Watershed Association Linda Vigil, Stenographer

2. APPROVAL OF THE AGENDA

Ms. McDonald will permanently change the order of the Agenda starting with the next agenda. Matters from the Commissioners will be moved before Information/Discussion/Action.

<u>MOTION:</u> Ms. Hansen moved to approve the agenda as presented with a second from Ms. Doremus which passed by voice vote.

3. APPROVAL OF THE MINUTES FROM July 14, 2016 and August 11, 2016

<u>MOTION:</u> Ms Doremus moved to approve the minutes of July 14, 2016 as presented with a second from Ms. Isaacson which passed by voice vote.

<u>MOTION:</u> Mr. Jacobi moved to approve the minutes of August 11, 2016 as presented with a second from Ms. Hansen which passed by voice vote.

4. COMMUNICATION FROM OTHER AGENCIES/COMMITTEES

There as not any communication form other Agencies or Committees.

5. INFORMATION/DISCUSSION/ACTION

A.) Project Update Review

Ms. McDonald presented the spreadsheet with the quarterly updates. (See Exhibit A)

Current Projects:

- Construction on Camino Alire, has started early.
- Arroyo Chamisos will be starting in October.
- Arroyo de los Pinos is in the final stages of planning. The final plans should be complete by next week.
- Drainage Repair Cross Vane at Calle Nopal will start late September or early October.
- Arroyo at Resolana Park is still working on grants for funding.

Ms. McDonald reported on projects that are complete. A discussion was held about several other projects not managed by Ms. McDonald but is relative to the River. Ms. McDonald discussed the way she reports projects to the Commission. Mr. Patorni asked about the reorganization of the City and the Departments that involve the River. Ms. McDonald is working on an RFP that will include that information.

Ms. McDonald reported the Stormwater Project will take up to two years to complete once it is approved.

A discussion was held about the St. Francis Gage. The grade control aspect of the project is complete.

A discussion was held about projects that require the flows to stop while under construction.

B.) River Parkway Benches and Bike Rack/PNM Power Up Grant Update

Ms. Baca-Tompson reported the grant was approved and the benches and bike racks have been ordered. A photo of the benches and bike racks and maps of their locations was shown. (See Exhibit B) Several of the benches will face the river. They will be installed by the end of the year.

Ms. Baca-Tompson discussed location 3 in Alto park will have the bench separate from the bike rack. The bench will be placed in front of the Rain Garden.

Mr. Otto discussed an event is being planned to dedicate the Rain Garden. He is working with Wells Fargo to coordinate it and will inform the Commission once it's planned.

Mr. Otto explained the projects were funded by a partnership between Wells Fargo and the US Fish and Wildlife. A small amount was also funded by the Santa Fe River Fund.

C.) Friends of Santa Fe River Walk Update

Ms. Hansen discussed the Friends of Santa Fe River walk that occurred in August. (See Exhibit C) A book was presented to Mr. Michael Smith and a walk of the river was done to see the progress of the removed invasive species of trees and grass.

Ms. McDonald discussed the project which was part of Parks. There are plans to reseed and replant in that area. There seems to be good feedback on the project.

D.) SF River Watershed Study Vegetation Mapping Report

Ms. McDonald presented a copy of the Vegetation Mapping Report (See Exhibit D). A copy was given to the species resiliency subcommittee to look at.

Ms. McDonald thought it was a great tool explains the soils and vegetation. It is an easy to read document and can be used for future goals of the Commission. Perhaps the idea to replant the vegetation reach by reach can be included in the goals. The Municipal Tree Board is already working on identifying the species of trees on the river.

Ms. McDonald explained it can be in the form of guidelines that can be outlined for other entities. The City would make it clear there are intentions of doing the work.

Mr. Bove discussed invasive grasses.

E.) Sub-Committee Breakout Session-Working Meeting

The Sub-Committee Breakout Session is not recorded.

- Outdoor Economy
- Promoting a Living River
- Watershed Revitalization
- Species Resiliency

6. MATTERS FROM COMMISSIONERS

Ms. McDonald discussed an email send by Chair Buchser (See Exhibit E). In the email the Chair asked the Commission to send 10-15 ideas to be discussed at the next meeting. Also a community meeting can be planned for May.

Ms. Hansen briefly discussed the way another committee she was a member of worked on a similar brain storming session.

Mr. Patorni stated there needs to be clarification of the River and the Watershed.

Ms. McDonald suggested the members create a list of as many ideas as they can and email them she will then combine and merge the ideas to one list. She will also suggest the idea Ms. Hansen had.

Ms. McDonald discussed the way the Water Conservation Committee ranked items.

Ms. McDonald briefly mentioned the small event for Mr. Smith (Friends of the Santa Fe River) went well although it was done with short notice. The Mayor would be happy to participate in the future. A brief discussion was held about how these events will be sent to the Commissioners ahead of time so they may attend.

Mr. Patorni would like a list of all other Committees and Commissions within the City. Ms. McDonald will help walk him though how to find them online.

Ms. McDonald discussed the form that was sent out from the City Clerk that releases telephone and contact information.

<u>MOTION:</u> Ms. Hansen moved to allow Ms. McDonald to share name, position on the committee, addresses and phone numbers and length of terms to send out with a second from Mr. Patorni which passed by voice vote.

A discussion was held about the staff list on the web for each City Department.

7. MATTERS FROM STAFF

There no matters to report from staff.

8. CITIZEN'S COMMUNICATION FROM THE FLOOR

Mr. Otto announced an appreciation picnic for volunteers of the River and Watershed will be on Saturday Sept. 24 from 11:00 am to 1:00 p.m. at the Water History Museum. The Commission is welcome to attend and perhaps speak on the history of the River. The building has been remodeled.

Mr. Bove discussed the Water History Museum and the History of the Power Plant Building. He would like to see old pictures of the River put up. Ms. McDonald will contact the proper department and boards.

Mr. Patorni briefly discussed the possibility of raising money or getting funds from the State Legislature.

9. ADJOURN

<u>MOTION:</u> Mr. Jacobi moved to adjourn the meeting at 7:22 p.m. with a second from Mr. Pierpont which passed by voice vote.

SIGNATURES

John Ruchser Chair

Linda Vigil, Stenographer

Public Works/Engineering Division

_
┖
5
Q
ē
œ
2
⊒
±
Sta
:=
Ś
ŭ
ಕ
Ψ
•=
Ç
=
Δ.

The state of the s				から ない はん から はまれる	
	Project Title	Project Manager	Contractor	Fund:Type(g)	
	SF River Small Drainage Repair - at Camino Alire		Constitution of the consti		
	Viela	Brian D / Melissa M	Garnett Fleming	2014 CIP Bond	Pecember 2014
MUB			1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
C		Melissa M	190	2014 CIP Bond	December 2015
	Construing Fosineering	Melissa M	SMA	2014 CIP Bond	
3 9	Construction Management	Melissa M	SFERB	2014 CIP Bond	
4					
		Bran D/ MAM	SEE	2012 GO Band	November 2015
3 6	5	a Xolan			
		WellstaM	BMC	2812 GO Bond	· December 2015
		MeissaM	SFE	2012 GO Bond	June sante
200		Mel ssa M	SMA/Sites SW	2012/GO Bond	
5	Design			44	
	400		, and	2012 Grilland	
6	Construction	Melssa W	30,1	200 - 200 E	
	Construction Engineering/Management	MelissaM	8	ZULA SEPONIO	
SOOR	III Drainage Rebair Cross Vane (@Nopal)				
		Melissa M	Sites SW	2017.60.8904	
0	MUB				
	Louis Inc.	Melissa M	Raincatcher		
を できる とう は できる とう		Meilssa M	SE	2014 CIP Bond	
25.55		Leroy P/Melissa M	NA	2012 GO Bond	
8 2					
4					
3				The second second	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Construction Engineering/Management				
		Control of the contro		100 miles (100 miles (× 200
412 A	SF River Small Drainage Repair - St. Francis Gauge		1	のでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、100mmでは、	
		Briano	Gannett Fleming	2014 CIP Bond	December 2014
		Melissa W	Lockwood	2014 CIP Bond	March 2016,
		Melista M	SWA	2014 CIP Bond	March 2016 -
=	Construction engineering management		Control of the contro		
	Arroyo Assessment		SFWA	2012 GO Bond	November 2015
	Professional Service Contract	(Michigae IV)			38 Sept. 14 Sept. 15

EXHIBIT

08/30/2016

City of Santa Fe, New Mexico Mexico Memory City of Santa Fe, New Mexico

DATE: July 27, 2016

APPROVED:

David Rasch, Historic Design Review Supervisor

VIA:

Leroy Pacheco, PE - River, Watershed & Trails Section Supervisor

FROM:

Melissa A. McDonald, RLA, River & Watershed Coordinator

ITEM/ISSUE:

PNM PowerUp Grant for benches and bike backs with in the Santa Fe River corridor

BACKGROUND/SUMMARY:

In March 2016 the Santa Fe Watershed Association with assistance from the City of Santa Fe 's River Commission applied for a grant to secure five benches and bike racks to be placed within the Santa Fe River corridor. This would allow for great enjoyment for the trail users. We were award the grant in May 2016.


The benches and bike racks specified were selected to match in style & color to the existing benches and bikes currently placed along the river specifically in the Parque del Rio park sites. Please see the attached packet with location maps, photos of existing benches, specification for new benches/bike racks, photo of proposed footing, and copy of Parque del Rio specification sheet.

The City of Santa Fe's Parks Department has agreed to install the benches and bike racks as part of the grant. The footings will be class "A" concrete with a minimum of 3,000 PSI compressive strength. Specification will match existing bench and bike rack installations per city standards.


RECOMMENDED ACTION: We are seeking approval of two bench & bike rack placement located within the historic district.


Attachments:

- Location Maps 1 5 with photo placement
- Bench/Bike Rack specifications
- El Parque del Rio Improvements specification sheet
- Estimate detailing type, color, quantity and cost









SCH Enterprises, LLC

79405 HWY 111, Ste. 9PMB 466, La Quinta, CA 92253

www.SCHenterprisesLLC.org operations@schenterprisesIlc.org 503-364-1353

PRODUCT SUBMITTAL INVERTED U SERIES – BIKE RACK

MODEL

	Model #	Description	Size
	Model #		21" W X 36" H
	SU 15	1.5" I.D. Schedule 40 pipe	
~		2" I.D. Schedule 40 pipe	21" W X 36" H
	SU 20		
	WU 15	1.5" I.D. Schedule 40 pipe	30" W X 36" H
لسا	AAO TO	-	30" W X 36" H
П	WILL 20	2" I.D. Schedule 40 pipe	20 M V 20 II

MOUNTING OPTIONS

- ☐ Embedded mount; E
- ☐ Flanged Surface Mounted; F

FINISH OPTIONS


- ☐ Hot-dipped galvanized; G
- ☐ polyester powder coated; P
- ☐ Thermoplastic, T
- ☐ Stainless steel, 304 with a #4 satin finish; SS

OPTIONAL

☐ CROSS BAR; CB



EXAMPLE: WU 20-F-SS



EXAMPLE: SU 20-E-P

SECTION 02870 - SITE FURNISHINGS

PART 1 - GENERAL

1.01 RELATED DOCUMENTS

A. Drawings and general provisions of the Contract, including General and Supplementary Conditions and other Division 01 Specification Sections, apply to this Section.

1.02 SUMMARY

- A. Drawings and general provisions of the Contract, including General and Supplementary Conditions and Division-1 Specifications sections, apply to work of this section.
- B The work of this section includes furnishing all site furnishings in accordance with the specifications and drawings contained in the Contract Documents. This shall include all labor, equipment and performance of operations including subsurface footings, connections, and cleanup of planting areas as specified herein.
- C. Related Work:
 - 1. Division 3, Section 03100 Concrete

1.03 REFERENCES

- A. ACI American Concrete Institute Manual of Concrete Practice
- B. ASTM American Society for Testing and Materials
- C. SPIS Society of Plastic Industry Standards

1.04 SUBMITTALS

- A. Product Data: Manufacturer's current printed specifications and catalogue cuts of the following:
 - 1. Park Bench Backed
 - 2. Picnic Table
 - 3. Picnic Bench Backless
 - 4. Trash Receptacle
 - 5. Bicycle Rack
 - 6. Dogi-pots
- B. Samples: Color and finish for each type of furnishing.

1.05 DELIVERY, STORAGE, AND HANDLING

A. Packaging and Labeling: Furnish materials in manufacturer's unopened, original packaging, bearing original labels showing quantity, description and name of manufacturer. Verify that materials and components are adequately padded and securely bound in such a manner that no damage occurs to the product during delivery and unloading at the site.

Picnic Table Backless Bench (For use with Picnic Tables)

- Manufacturer: Creative Pipe 1.
- Model No: Granada Backless Bench, BCBLB-FB-6(or 4')-SM-T, 6' long or 4' long see plans. 2.
- Material: Steel 3.
- Finish: Thermal-plastic powder coat 4.
- Color: Brown range. Submit metal color samples for Owner select. 5.

Trash Receptacle

- 1. Manufacturer: Creative Pipe
- Model No: BCT-FBS-32-F-T-LD 2.
- Material: Steel 3.
- Finish: Thermal-plastic powder coat
- Color: Brown range. Submit metal color samples for Owner select. 5.
- Accessory: Order matching trash container insert by the manufacturer. 6.



Bicycle Rack

- 1. Manufacturer: Creative Pipe
- Model No: SU-20-F-T 2.
- Material: Steel pipe 3.
- Finish: Thermal-plastic powder coat 4.
- Color: Brown range. Submit metal color samples for Owner select. 5.

F. Doggy Bag Dispenser

- Manufacturer: Dogi-pot 1.
- Model No: 2.
- Material: Steel 3.
- Finish: Thermal-plastic powder coat 4.
- Color: Brown range. Submit metal color samples for Owner select.

Drinking Fountain

- Manufacturer: Most Dependable Fountain 1.
- Model No: 2002 Historic Fountain 2.
- Material: Steel 3.
- Finish: Thermal-plastic powder coat
- Color: Brown range. Submit metal color samples for Owner select.

MANUFACTURER'S INFORMATION 2.03

Submit copy of manufacturer's information on site furnishings in Operations and Maintenance manual for project.

PROTECTION 2.04

Do not remove protective wrappings from furnishings until instructed by Landscape Architect. A.

SCH ENTERPRISES LLC

79405 Hwy 111 Ste 9 PMB 466 La Quinta, CA 92253 (503) 364-1353, CA US (760) 340-5555 accounting@schenterprisesllc.org, manager@schenterprisesllc.org, operations@schenterprisesllc.org



ADDRESS

Santa Fe, City of-NM


SHIP TO

Santa Fe, City of-NM Raquel 505-920-4166

ESTIMATE 1743

DATE 07/25/2016


ACTIVITY	QTY	BATE	AMOUNT
BENCHES:BCBB-FB-6-SM-T Banico Series Backed Bench, Flat Bar Slats, 6' in length, Flanged Surface Mount, Thermoplastic Powder Coat Finish	5	845.00	4,225.00
INVERTED U:SU 20-F-T Inverted "U" Series Bike Rack, 2 Bike Capacity, Flanged Surface Mounted, Thermoplastic Powder Coated Finish	5	155.00	775.00
FLGMT This price is for a Flanged Surface Mount Product to be bolted to concrete.	1	0.00	0.00
THERMO The above price is for a single coat standard color Thermoplastic powder coated finish - BROWN	1	0.00	0.00
PRICE LOCK SCH Enterprises, LLC will hold the above listed price for thirty (30) days from date of this estimate.	1	0.00	0.00
SHIPPING Shipping Charge to Santa Fe, NM for five (5) benches and five (5) bike racks	1	1,598.00	1,598.00
FUEL Due to the current fluctuation in fuel surcharges, quoted freight is only an estimate. Actual freight charge will be billed at time of shipment.		0.00	0.00
ED Please note if residential or inside delivery is required, there may be a separate freight charge which will be billed to you after delivery. Charge will be a minimum of \$60.00.	1	0.00	0.00




EXHIBIT











MCDONALD, MELISSA A.

From:

ROSS, MATT

Sent:

Tuesday, August 23, 2016 3:46 PM

To:

ROSS, MATT

Subject:

RELEASE: City Hosts River Walk in Appreciation of Friends of the SF River Work



City to host walk along Santa Fe River Parkway to say thanks to the Friends of Santa Fe River and Michael Smith

FOR IMMEDIATE RELEASE

Date: August 23, 2016

Contacts:

Melissa McDonald, River and Watershed Coordinator (505) 955-

6840, mamcdonald@santafenm.gov

Matt Ross, Public Information/Multi-Media Administrator Officer, (505) 955-

6045, mross@santafenm.gov

Santa Fe, NM - Michael Smith, who leads the non-profit *Friends of the Santa Fe River* will wrap up three weeks of river riparian invasive-species removal at the end of this week. The work was done in cooperation with the City of Santa Fe Parks and Recreation Department, Fire Department, and the River, Watershed and Trails Section of the Public Works Department.

Most significantly, Michael's passion for the work succeeded in inspiring the cooperation and financial participation of a number of donors who support the work to remove invasive Siberian elms from the river parkway. This public/private partnership is founded upon a cooperative agreement that was approved by the city in 2015. We look forward to continued cooperation between the City of Santa Fe and *Friends of the Santa Fe River* in the years ahead.

In appreciation for all of the work accomplished in the last few weeks, the City of Santa Fe will host a public walk of the site with Michael Smith. The public is invited to attend the walk and learn more about the work being done in the riparian zone of the river. The short walk will commence at 11 AM on Thursday, August 25th, 2016 and will take no more than 30-minutes. Those interested should meet at the park tables along the Santa Fe River just south of El Alamo Street.

FRIENDS OF THE SANTA FE RIVER

http://www.friendsofthesantaferiver.org/




Santa Fe River Watershed Study Vegetation Mapping Report




1. Introduction

In December 2007, a Report on Existing Conditions and Potential Natural Resources Restoration Projects (Blue Earth) was completed as part of the Santa Fe Watershed Study. This report covered existing ecological conditions in the original study reach, which was from Camino Alire Rd. to where the river passes under NM 599 (approximately 8.4). In 2008, the study area was expanded to include additional portions of the river both upstream and downstream of the original study area. This included an additional 1.6 miles below NM 599, an additional 4.5 miles above Camino Alire Rd, as well as four arroyo tributaries: Arroyo Mascaras (1.6 miles), Arroyo de la Piedra (2.6 miles), Arroyo Ranchito (1.8 miles), and Arroyo Barranca (1.9 miles), for a total of 14 additional miles (Figure 1). The original study area is show in Figure 2.

In 2009-2011, an inventory of existing vegetation of these additional areas was performed. The remainder of this document will discuss that mapping, how it integrates with the previous vegetation inventory, and the overall results.





2. Channel Morphology

2.1 Classification of Stream Segments in the additional Study Area
The Blue Earth report described causes for channel incision in the Santa Fe River. The additional portions of the study contain similar incision issues.

When the Corps performed vegetation mapping on the additional areas, some channel morphology measurements were also taken. Downstream of 599, the channel is fairly wide until it reaches the treatment plant. At this point the channel becomes more constricted just above where 599 crosses the river.

Where the river runs through the center of town, the channel is very constricted and has rock walls defining its edges through much of this stretch (Figure 3). Development including homes and businesses line the river on both sides throughout this area.




Figure 3. Portion of the Santa Fe River along West Alameda Street

The upper portion of the Santa Fe River includes an Audubon Wildlife Preserve Area just downstream of Upper Canyon Dam. In this area, the channel is wider and more natural but still bounded by large individual lots on either side (Figure 4).



Figure 4. West bank of the Santa Fe River downstream of Upper Canyon Dam The Arroyo Mascaras runs northeast from the Santa Fe River starting at North Saint Francis Drive. It runs along the south side of Paseo del Peralta and bends to the north before becoming Arroyo Barranca. The channel is flat and open (Figure 5) until it meets up with Arroyo Barranca at the Fort Marcy Park where it is heavily used for recreation (Figure 6).



Figure 5. Arroyo Mascaras




Figure 6. Arroyo Mascaras where it becomes Arroyo Barranca Arroyo Barranca begins just above Camino Encantado and runs down to where it meets/becomes Arroyo Mascaras above the Fort March Park at Murales Road. The channel is wide and flat. Development is comprised of large open lots that are spread out on either side of the river with limited structures directly adjacent to the Arroyo (Figure 7).



Figure 7. Arroyo Barranca

Arroyo de la Piedra and Arroyo Ranchito were not visited due to time and budget constraints of the study.

The Santa Fe River downstream of 599 includes a portion near the treatment plant. In this area, the channel is more constricted but contains lush native riparian vegetation along the banks (Figure 8). Between the treatment plant and 599, the channel is more open with sparse vegetation (Figure 9).



Figure 8. Santa Fe River channel downstream of the treatment plant




Figure 9. Santa Fe River channel between 599 and the treatment plant

3. Plant Communities

Plant communities in the study area were mapped using the community-structure (C-S) classification scheme developed by Hink and Ohmart (1984). This classification combines identification of community dominants in the tree and shrub strata with the structural character of the stand being delineated, where structural character is defined as the variation in foliage density with height above the ground surface. From 2009-2011, the Corps mapped the vegetation within the additional study areas using the same methodology employed by Blue Earth for the original study area. The Corps completed mapping along the areas added to the mainstem of the Santa Fe River as well as the Arroyo Mescaras and Arroyo Barranca. Mapping was not completed on Arroyo de la Piedra or Arroyo Ranchito.

Six structure types are used in the classification. These range from structure type I, characterized by an overstory canopy provided by mature trees (i.e. 50 to 60 feet tall) and understory foliage to type VI, characterized by sparse herbaceous and shrubby vegetation. A seventh structure type, X, was added to the classification to describe lacking woody dominants and with foliage restricted to three feet above the ground and lower. This structure type was used in the original analysis by Blue Earth. Two miscellaneous cover classes were used for area lacking woody vegetation. The code BARE was used for areas with sparse herbaceous cover and greater than 70 percent bare ground. The code HERB was used for areas dominated by herbaceous plants. The only difference was that the Corps did not use the 'Bare X' category employed by Blue Earth. This was used in the original study area "to describe lacking woody dominants and with foliage restricted to three feet above the ground and lower" (Blue Earth, 2007). Instead, the Corps

utilized the OP (open) code when required (when the majority (~70-100%) of the polygon contained no vegetation and also utilized the Type VI structural code for 'very young, low, and/or sparse stands, the majority of which is between 0-5 feet in height' per the Hink and Ohmart protocol. There were only two cases where this was required, on the south end of the Santa Fe River between 599 and the new southern end of the study area.

Plant community types were classified using a combination of 16 codes for dominant or codominant species or cover type (3 codes were added for the new study area). Nine of the codes were for native woody or suffrutescent plant species: C for cottonwood (including Rio Grande, narrowleaf, and lance-leaf), CW for coyote willow, GW for Goodding's willow, J for one-seed juniper, LO for New Mexico locust, RB for rubber rabbitbrush, P for pinon pine, MM for mountain mahogany, and BE for Box elder (Table 1). Another five codes were for non-native woody plants: HL for honey-locust, RO for Russian olive, SC for saltcedar, SE for Siberian elm, and TH for tree-of-heaven (Table 1).

Dominant herbaceous species in areas delineated as HERB included hairy golden-aster (Heterotheca villosa), horseweed (Conyza canadensis), smooth oxeye (Heliopsis helianthoides), sand-daisy (Dieteria canescens), rough cocklebur (Xanthium strumarium), prickly lettuce (Lactuca serriola), bur ragweed (Ambrosia acanthicarpa), Russian-thistle (Salsola tragus), white sweet-clover (Melilotus albus), sorrel wild-buckwheat (Eriogonum polycladon), cañaigre (Rumex hymenosepalus), Canada wildrye (Elymus canadensis), Indian ricegrass (Achnatherum hymenoides), cheatgrass (Bromus tectorum), Carolina lovegrass (Eragrostis pectinacea var. pectinacea), and foxtail barley (Hordeum jubatum).

CODÉ	SPECIES/COVER 1		
BARE	Mostly bare ground with scattered herbaceous plants		
HERB	Herbaceous vegetation		
	<u> </u>		
С	Rio Grande, narrowleaf, and/or lance-leaf cottonwood; lance-leaf is a hybrid between Rio Grande and narrowleaf (<i>Populus deltoides wislizenii</i> , <i>P. angustifolia</i> , and/or <i>P. x acuminata</i>)		
CW			
GW	Goodding's willow (Salix gooddingii)		
	one-seed juniper (Juniperus monosperma)		
LO	New Mexico locust (Robinia neomexicana)		
RB	rubber rabbitbrush (Ericameria nauseosa)		
Р	Pinon pine (Pinus edulis)		
ММ	Mountain mahogany (Cercocarpus montanus)		
BE	Box elder (Acer negundo)		
HL*	honey-locust (Gleditsia triacanthos)		
RO*	Russian olive (Elaeagnus angustifolia)		
SC*	saltcedar (Tamarix chinensis)		
SE*	Siberian elm (<i>Ulmus pumila</i>)		
TH*	tree-of-heaven (Ailanthus altissima)		

Table 1. Species/cover codes used in describing plant community types in the study area. Scientific and common names follow Allred (2006). Those species marked with an asterisk (*) are non-native.

The lower end of the Santa Fe River from Highway 599 down to just below the treatment plant was mapped in 2009-2010. The area between Highway 599 down to the treatment plant is similar to the river above Highway 599 with an open wide channel and sparse vegetation comprised of more upland species such as sand sage and juniper with some riparian species mixed in (Figure 10). The area just below the treatment plant, however, contains a diverse mix of riparian vegetation due to the continual water source from the treatment plant. This area is dominated by cottonwood, coyote willow and tree willow with some Siberian elm and Russian olive mixed in (Figure 11). The understory is also very thick in much of this area creating Type I and Type III stands. The habitat in this area is a good example of what potential restored habitat should mimic.



Figure 10. Santa Fe River between 599 and the treatment plant



Figure 11. Vegetation along the Santa Fe River downstream of the treatment plant

The upper end of the Santa Fe River that was not in the original study area was mapped in 2010-2011. This upper portion of the river was mapped from the Audubon Preserve Area just below the Upper Canyon Dam down to the original study area boundary (St. Francis Drive). The upper portion of this stretch starting at the Preserve Area contained more riparian and montane species with cottonwood, box elder, coyote willow and tree willow being dominant species (Figure 12). Box elder (Acer negundo) was added to the C-S list due to its presence in this area. Some additional woody species that were detected in this area include: coral berry (Symphoricarpos orbiculatus), Wood's rose (Rosa woodsii), spruce (Pinaceae), and currant (Ribes spp.).




Figure 12. Vegetation downstream of Upper Canyon Dam

Arroyo Mascaras also runs northeast from the Santa Fe River but to the east of Arroyo Barranca and Arroyo Ranchito. Plant C-S types in this arroyo were dominated by the following native woody species: Siberian elm, rabbitbrush, and some small patches of cottonwood and coyote willow (Figure 13).



Figure 13. Vegetation along Arroyo Mascaras

Arroyo Barranca runs northeast from the Santa Fe River and ends just above Camino Encantado. Pinon pine (*Pinus edulis*) and mountain mahogany (*Cercocarpus montanus*) were added to the C-S list due to their presence in this arroyo. Plant C-S types in this arroyo were dominated by the following native woody species: juniper, pinon pine, mountain mahogany, and rabbitbrush (Figure 14). The majority of stands were Type IV structure.







Figure 14. Vegetation along Arroyo Barranca


Major factors influencing the current condition of riparian vegetation in the study area are 1) significantly reduced surface water flow; 2) loss of the shallow alluvial aquifer; 3) massive bed degradation throughout the reach initiated in the mid 1970s by removal or lowering of grade control structures; and 4) scouring peak flows associated with storm-water runoff that are now contained within a narrow, entrenched valley throughout most of the study area. Because of these factors, riparian vegetation is sparse throughout the study area and where it is found it is typically characterized by early successional stages.


Vegetation maps of the entire study area are provided as a separate 11x17 map book document.




























































































Report on Existing Environmental Conditions and Potential Natural Resources Restoration Projects

5 December 2007

Prepared for U.S. Army Corps of Engineers, Albuquerque District Delivery Order 0014, Contract No. DACW47-03-D-005



Prepared by
John S. Pittenger and Karen A. Yori
Blue Barth Ecological Consultants, Inc.
1345 Pacheco Street
Santa Fe, New Mexico 87505



U.S. Army Corps of Engineers, Albuquerque District

3.2.2 Santa Fe River Channel Morphology	, ,
man at the other and Malan	~
2 C. Carinana amin Desironment	_
3.6 Land Use	-
.0 POTENTIAL NATURAL RESOURCES RESTORATION PROJECTS	3
a man at the standard Destruction?	,,
4.3. Producedon Confered Objectives	•
a n a tradition - Destruction Goal	•
and the state of t	,,
a man and a management of Projects	•
And the Bloom Danger	-
A D. Co	• •
and beat Similar Manufation	•
4.3.4 Community Involvement	69
5.0 PREPARATION, COORDINATION, AND CONSULTATION	70
5.1 Preparation	70
5.2 Coordination and Consultation	70
6.0 REFERENCES	71
APPENDIX A Animal Species Potentially Occurring in Riperian or Aquetic Habitata in Santa Fe Cour	ij
APPENDIX B Guidelines for Ripstism Planting	84
APPENDIA D CHEGORIO LOS REPOSIDO LOS DELINOS	

LIST OF TABLES

The state of the s	í
Table 1. Resgen stream type classification of channel segments in the study area	ı
m at a County amount atmosphere in the study area.	
The state of the s	•
m 14 4 Commission was added used in describing plant community types in the study area	•
make a Community of the	•
	•
1 Sole 7. Companions of the second of Minney or non-Lating residents	į
Table 8. Proportion of Hispanic or Latino and non-Hispanic or non-Latino residents	¢
TO U.S. D. Communicate of colocted income data	•
Table 10. Acreage by land ownership in the study area	١

U.S. Army Corps of En

TABLE OF CONTENTS

.0	INTRODUCTION	÷
	1.1 The Study Area 1.2 Past Restoration Projects That Currently Influence Existing Conditions	•
	EXISTING ENVIRONMENTAL SETTING	3
2.0		
	2.1 Climate 2.2 Physiography and Geology	4
	2.2 Physiography and Geology	9
	2.2 Physiography and credogy 2.3 Groundwater	9
	2.3 Groundwater 2.3.1 Tesuque Formation	10
	2.3.1 Testique Permanon 2.3.2 Terrace Deposits and Recent Alkuvium	ii
	2.3.2 Terrace Deposits and Recent Attavalant	ii
	2.4 Surface Water 2.4.1 Hydrology	ï
	2.4.2 Santa Fe River Channel Morphotogy 2.4.2.1 Causes of Channel Incision and Destabilization	16
	2.4.2.2 Classification of Stream Segments in the Study Area	24
	2.4.2.2 Classification of Structures in the Study Area 2.4.2.3 Grade Control Structures in the Study Area	24
	2.4.2.3 Grade Control Structures in the Solidy Acce. 2.4.3 Water Quality	27
	2.4.3 Water Quality 2.5 Soils	14
		7.
	2.10.1 Camino Carlos Rael to CR 62 (Santa Fe County)	-
	2.10.4 Cottonwood Drive to NM 599 (Santa Fe County)	31
3.	0 FUTURE CONDITIONS WITHOUT RESTORATION	
-		
	3.2.1 Hydrotogy and Water Quality	•

Senta Fe River Study

Page

U.S. Army Corps of Engineers, Albuquerque Distr

LIST OF FIGURES

	~
igure 1. Location of the Santa Fe River study area	2
Figure 38. Rabbitbrush (RB VI) regenerous on a thoughtain the area below Camino Carlos Rasi	40
Figure 37. Simili or Separate duri asses)	

Figure 40.	Remnant hydric soil indicators in a cut bank below CR 62 at Agua Fria 4
Figure 41.	Land use and ownership features, Camino Alire to Camino Carlos Rasl
Figure 42.	Land use and ownership features, Camino Carlos Rael to below Siler Road
Figure 43.	Lend use and ownership features, below Siler Road to below CR 62
Figure 44.	Land use and ownership features, below CR 62 to Cottonwood Drive
Figure 45.	Land use and ownership features, Cottonwood Drive to NM 599
	Waste piles spilling into the river channel
Figure 47.	Unauthorized grading plugged the restored meander constructed in 2000
	View upstream of the restored meander now isolated from the river
Figure 49.	Grade control structure GCS-7, downstream from Camino Carlos Rael
	Coyote willow establishment on a point bar
	Storm drain outfall SW-12 at Siler Read
Figure 52.	Stand of narrowleaf cuttonwood at structure SW-5
	Conceptual design of a storm-water detention basin
	Storm-water detention pond near the NM 599 bridge
D: 66	Conserved design for significant and assert a server and asset at the server

LIST OF PLATES

Plate 1. Plant community-structure types in the study area	, station 0+00 to 86+00
------------------------------------------------------------	-------------------------

- Plate 2. Plant community-structure types in the study area, station 86+00 to 164+00

- That 3. Plant community-statucture types in the study area, station 164-00 to 246-00 Plate 3. Plant community-statucture types in the study area, station 164-00 to 246-00 Plate 4. Plant community-structure types in the study area, station 246-00 to 330-00 Plate 5. Plant community-structure types in the study area, station 330-00 of 378-02 Plate 6. Rosgen classification of atream segments and locations of grade control structure. outfalls in the study area

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District



Location of the Santa Pe River study area (blue line) in northcentral Santa Fe County, New Mexico. Major tributuries to the river in the study area are labeled. The portion of the Santa Fe River watershed on the mapped area is indicated by yellow shading.

1.0 INTRODUCTION

This report describes existing environmental conditions and potential natural resources restoration projects for a segment of the Santa Fe River in and near the City of Santa Fe, River Mexico (Figure 1). The Santa Fe River was declared America's most-endangeredriver in 2007 by American Figure (2007). by American Rivers (2007).

The report is part of a larger General investigation. Watershed Study by the U.S. Army Cosps of Engineers, Albuquerque District, in cooperation with the City of Santa Fe and Santa Fe County. The purpose of the study is to 1) identify alternatives for restoration of the structure and fametion of the Santa Fe River and 2) determine the federal internet in watershed planning and water resources mesagement in the study area.

1.1 The Study Area

The Sania Fe River study area is tocated in noeth-central Sania Fe County, New Mexico (Figure 1). The study area begins at the Camino Alire bridge crossing within Sania Fe city limits and extende downstream 7.16 miles to the NAM. Highway 599 bridge crossing. For the purposes of this report, the study arms in defined as the area coologically influenced by the river. This was interpreted in the field as the active river classnet, established and associat floodplain areas, and low termoes potentially subject to flooding under relatively commonplace flows. Field mapping and photo-interpretation were employed to delineate the area of river influence, which ended up encompassing about 83 acres. The boundary of the area of river influence, which ended up encompassing shout 83 acres. The boundary of the area of river influence, is shown on figures depicting soil types and plant communities in the study area (cf. sections 2.4 and 2.8). The Santa Pe River study area is located in north

1.2 Past Restoration Projects That Currently Influence **Existing Conditions**

Two major grade control structures in the study area, while not constructed specifically for the purposes of ecological restoration, play a major role in arrowing bed degradation and establishing channal equilibrium. These functions are critical with respect to development of ripasian registrion and other scological attributes in the study area. The County Road 62 (CR 62) crossing was constructed in 1994 and the San Yeldro crossing was constructed in 1999. Both structures were designed by the U.S. Army Corps of Bagineers, Albuquerup District. designed by the U.S.. Albuquerque District.

A 1.5-mile restoration project on State Trust lands at the downstream end of the project area was implemented in 1999-2000. This project involved removal of a road-fill creasing, construction of a meander at the crossing site, installation of several root-wad revolucests and extensive planning of riparian vegetation to premote nearowing of the channel and meander development.

Another restoration project was conducted on the reach between the Sm Yeidro crossing and CR 62 in 2005. This project involved excavation steep, ereding basics and reconstrucing them to a more gentle slope, channel reconfiguration, installation of a grade control and energy dissipation structure, and siperion plentings.

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

2.0 EXISTING ENVIRONMENTAL SETTING

2.1 Climate

Climate of the study area is dry semi-strid (Köppen-Geiger classification BSIs; Kottek et al., 2006). Precipitation is concentrated in the summer when southeasterly circulation of air meases brings moist air up from the Oulf of Mexico. Storag surface basting and orographic lifting causes these air meases to rise, resulting in condensation of moistners and rainfall. In contrast, winter precipitation derives primarily from Pacific Ocean storms that are strong enough to persist inland to New Mexico. Such frontal systems west to past across the state.

Average maximum summer temperatures are around 80°F, with associated average low temperatures in the mid-50s. Winter average high temperatures are in the low to mid-40s with lows typically near 20°F (Figure 2). Precipitation is concentrated in the summer months. Byspotranspiration increases markedly in April. average must be considered in April and stays high through the growing concon (Figure 2). Average annual precipitation in the study area is about 13.7 inches but there is considerable variation from year to year. The lowest renerable variation from year to year. The lowest recorded annual precipitation, 5.03 inches, was in 1917 while the highest annual precipitation (21.75 inches) on record was in 1881 (Figure 3).




Figure 2. Climate characteristics for the Santa Fe area, 1867 to 2006. Data for 1867 to 1972 are from the to Fe station. Santa Fe station (298072), 1972 to 2006 data are from the Santa Fe 2 station (298085; Wastern Regional Climate Center, 2007). Class A pan evaporation data for 1867 to 1972 are from Oregon Climate Service (2007). Pan evaporation data for 1971 to 2006 were not available.

Figure 3.
Precipitation and drought record for the Santa Pe area, 1867 to 2006. Precipitation data are from stations 298072 and 298085 298072 and 298085 (Western Regional Climate Center, 2007) as described in the caption for Figure 2. Palmer drought severity index data are annual averages. Monthly index data are imm National Climatic Data Center (2007).

Since climate data in the Santa Fe area began being recorded in the late 1800s, drought conditions (i.e. a Palarner drought index of 4.0 or lower) were observed in the area around the turn of the 20th century, during the 1950s, from 1963 to 1964, in 1967, and from 2000 to 2004 (Figure 2). 1 New Mexico Climate Center, 2007).

2.2 Physiography and Geology

Located in the Southern Rocky Mountains province of the Rocky Mountain System physiographic region (Fennesma and Johason, 1946), the roudy area is situated on a broad plain or piedmost on the east side of a trough formed by the Rio Grande rift (Shrobs et al., 2005). The trough of the Rio Grande rift in the vicinity of


Santa Fe is bounded on the east by the Sangre de Cristo Mountains and on the west by the Sierra Nacimiento (Spiegot and Baldwin, 1963: 6). The portion of the Rio Grande rift valley including the study area is known as the Santa Fe Embayment.

Erosion of the pre-Tertiary rocks of the Sangre de Cristo Mountains during the late Oligoceae to Mioceae (ca. 33 to 23 million years ago, Read et al., 2000), initiated by upilf of the mountain sediments in the subsiding wallal., 2000), initiated by uplif of the mountain mass, resulted in extensive deposition of detrial sediments in the subsiding valley basin. These deposited sediments are known as the Tesuque Formation of the Santa Fe Group and they remaitos of the Santa Fe Lincop and they comprise the major water-bearing goologic formation in the Santa Fe zrea (Spiegel and Baldwig, 1963: 133, Lewis and West, 1995; Strobe et al., 2005; 11). The Tesuque Pormation comsists of light brown to red unconsolidated

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

Figure 4. Geologic map of the upstreamhalf of the study area, excepted from Read et al. (2000). The inset diagram of stratigraphic units was adapted from Figure 5 in Konig et al. (2002).



U.S. Army Corps of Engineers, Albuquerque District

nents and sandstone (Shrubs et al., 2005: The upper unit of the formation is asolidated to weakly consolidated sediments unconsolidated to weakly consolidated sediments varying from slightly outboly pebble gravel to sithy sand, while the lower unit is very fine to fine sand, sandstone, and sandy pebble conglornerste in tenses and tabular beds (Shroba et al., 2003: 16-17). The Tesuque Formation in the study area overtices Proterousic or Pennsylvanian rocks and is in excess of 3,700 feet thick (McAda and Wasiolck, 1988: 64).

Wasiolek, 1988: 64).

Aggradation of the Tesuque Formation ceased about eight million years ago (Konig et al., 2001). The Tesuque Formation is overlain on the piedmont surface north and south of the Santa Fe River by another depositional feature, the Ancha Formation (Konig et al., 2002). The Ancha Formation consists of light beownish grave to brown sity sand to pebbly sand with gravel and was deposited by atreams draining the touthern Sanger de Cristo Mounhains during the late Pilocene to sarly Pleistocene time (Konig et al., 2002; Shroba et al., 2005). The Ancha Formation years ago (Konig et al., 2002; 203), probably because of inceision of the Santa Fe River through the relatively resistant rocks of the Corror del Riscolantic field at La Bajada and its subsequent drainage of the Santa Fe cenbayment of the Sportation ranges in thickness from about 10 to 200 feet (Shrobs et al., 2005). This formation has been redord and reworked in the study area by the Santa Fe River. the Senta Fe River


A third stratigraphic unit was deposited on top of the Aucha Formation in the early Pleistroceae (Kong et al., 2002; Shrobs et al., 2005). This unit, mapped as shewtwash deposits by Shrobs and others (2005). This unit seny reach thicknesses of up to about 15 feet in the vicinity of the study area (Shrobs et al., 2005; 7).

Surficial geology in the study area consists of modorn alluvium associated with the active channel of the Santa Fe River and finaling terrance deposits actionasts that represent floodplains formed at higher base levels by the annostral Santa Fe River (Figures 4 and 5). Actificial fills associated with old gravel pits and landfills are common in the lower half of the study area, between County Road 62 (CR 52) and N.M. Highway 599 (NM 599; Figure 5).

Exposures of the Tseuque Formatine are evident along the vertical banks of the incised river channel throughout the stady area (Figure 6). Outcrops of consolidated Tseuque Formation analestone in the bed of the river channel were uncountered sporadically in theriver channel from encountered sport-dicatly in the river Common treat the sold stream gaps downstream from Carotino Alire to below the CR 62 crossing (Figure 7). Outcopes of Tenqua Formation in the chamal-bed were absent in the remainder of the study area, from about 2,000 feet downstream from the CR 62 crossing to the NM 599 bridge.

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque Di



coted Grow Shrobn et al. (2005). The inset diagram of stratigraphic units was adapted from Pigure 5 in Konig et al. (2002).




Figure 6. Exposure of the Tesuque Formation below Quatemany alluvium on the left (south) bank of the Santa Fe River upstream from Frenchy's Park, 15 June 2007 (photo by J. Pittenger).




Figure 7. Tesuque Formation outerop in the bed of the Santa Fe River at San Ysidro River Park June 2007 (photo by J. Pittenger).

U.S. Army Corps of Engineers, Albuquerque District



Pigure 8. View upstream of the Cieneguin Spring site at present-day Prenchy's Park, 26 June 2007. Arrow indicates outcropping of compact, and conglomeratic sitts of the Tesuque Formation (photo by J. Pittenger).

Groundwater may be present at the top of the Teasure Formation within about 1.5 miles of the front of the mountains but it generally much deeper at locations further out from the mountains (i.e. at depths greater than 50 feet, Lazanas and Drakes, 1995; Grant, 2602: 12), such as in the study area. Due to the complex, heterogeneous composition of the Tesuque Formation, depth to composition of the Teauque Formation, depth to groundwater may vary considerably from one location to the next (Lewis and West, 1995). The occurrence of shallow groundwater at the top of the Teauque equifor is difficult to predict because is associated with the presence of discontinuous paleochannels eroded into the top of the formation (Lazarus and Drakos, 1995).

ater from the Tesugue aquifer is hard due to high calcium and bicarbonate, which are the major chemical constituents in the water Hardness generally ranges from 121 to 232 parts

per million (ppm), total dissolved solids typically range from 90 to 500 ppm, and electrical conductivity is low and ranges from 150 to 800 micromiton per continuative (Spingel and Baldwin, 1963: 134; Lewin and West, 1995).

2.3.2 Terrace Deposits and Recent Alluvium

Terrace deposits and recent alluvium occur as thin Terrisco doposits and recent alluvism occur as thin (ca. five to 40 feet thick, Lezarus and Drakos, 1995) sharets of course, unconsolidated sediment overlying the Tesuque Formstion in the study area. Historically, the alluvium held a perched or semi-perched water table maintained by rainfall and infiltration of surface water in the Satats Fe River in locations where the underlying Tesuque Formation is relatively impremeable (Spiegel and Baldwin, 1963: 138-143). Where alluvium is

2.3 Groundwater

The following discussion presents basic information on groundwater characteristics that are relevant to ecological restoration in the study including depth to groundwater and adwater flow patterns in the study area. It is promise two process is the study area. It is not an exhaustive or comprehensive analysis of groundwater in the shudy area. The water-bearing goologic units in the shudy area, from oldest to youngest, are the Tesuque Formation, terrace deposits, and recent alluvium. Presence of water in necess alluvium is of most importance to restoration of ripation vegetation in the study area.

2.3.1 Tesuque Formation

Z.D.1 A Esuque Formation

The Tesuque Formation comprises a deep squiffer and, as discussed in section 2.2, is an important value source for the Santa Fe area. Recharge of the Tesuque Formation occurs at the mountain front where the formation outcrops, through allavium in ittram channels such as the Santa Fe River, and via areal recharge from percolation of precipitation through overtying sectiments (McAda and Wesieldel, 1938: 29-30). A ridge in the potentiomentic surface of the Tesuque squiffer under the Santa Fe River attest to the fact that midilitation of surface water flows and stabulow groundwater moving through allavium in the channel do recharge deep groundwater (Spiegal and Browster, 1963: 131; Lewiss and West, 1995). Croundwater flow in the Tesuque aquifer is from satt to west-outhwest in the study area (Lewis and West, 1995). and West 1995)

Groundwater in the Teruque Formation discharged naturally into the Santa Fe River in at least two locations in the study area as recently as the early 1960s. Seeps and springs discharged into the Santa Fe River at Cieneguita (present-day

location of Pranchy's Park; Pigure 1) and at Agua Pria (Spiegel and Baldwin, 1963: 132; Figure 1). The occurrence of these springs was described by Spiegel and Baldwin (1963:132) as follows:

The discharge (of groundwater from the Temopus Formation) at Clonoguita is probably the result of impedance of ground-water flow through the permeable rand section of the the permenous same section by the rue formation by compact red purposes sits which crops out in the el of the Santa Fe River just below the chammed of the Santa Fe Rhver just below the west limit of Santa Fe. The native growth of cottonwoods, the emergence of ground water, and the shallow water table to the Tesaque formation upstream — all indicate the existence of a partial barrier. Probably some ground water leaks through the barrier at ground water leafs through the barrier at depth, as well as flowing down the channel cut into it. Downstream from the barrier, the ground-water discharge probably returns to the Teauque formation, as the water slabe there is deep again. Similar conditions provail at Agas Pria, where also the ground water in a sandetone of the Teauque formation flowe across a barrier. The overflow emerges in the Santa Fa Biver channel became it is lowest possible overflow line in the a outhesis added).

The Clonegaita Spring (Figure 8) and Agua Fria Spring sites on the Sixths Fo River on longer have surface water except during periods of rainfall mooff, due to lowering of groundwater levels by groundwater withdrawal from municipal wells, reduced stream flow, and coession of irrigation. Currently, groundwater in the Tesuque Formation is well below the surface in the study area (McAda and Wasiotek, 1988: 23) and therefore is not likely to influence existing plant growth or ecological function.

Santa Fa River Study

U.S. Army Corps of Engineers, Albuquerque District

undertain by more permeable deposits of the Tesuque Formation, water infiltrates from the alluvium into the deeper aquifer (Spiegel and Baldwin, 1963; 141).

This althviral equifer was of sufficient volume that it discharged as springs and seeps at various locations along the river, most notably in the vicinity of Cinengs Street in downtown Santa Fe (Spiegel and Baldwin, 1963: 139). Similarly, several puebles were located along the river at cienages or marrity areas that were supported by a high water table in the alluvium (Spiegel and Baldwin, 1963: 19-3). These sites, including Los Palaccios, Pino, and Pueblo Quennado, were later settled by Spanish immigrants because of the water available for irrigation (Spiegel and Baldwin, 1963: 94). Many older homes sleng the Santa Fe River had shallow hand-day wells completed in the alluvial aquifer (Lazarus and Drakot, 1995).

Perched weter no longer occurs in the allavium and springs, sceps, and wells supplied by this stallow aquifer have gone dry since the early 1960s (Lezanus and Drakos, 1995). Loss of the allavial aquifer is attributeble to elimination of flow in acceptass and cessation of flood-irrigation, flow in acequies and cessation of flood-brigation, groundwater pumping, reduction of stream flow, and paving-over of recharge areas (Spiegal and Baldwin, 1963: 141-142; Lezarus and Drakos, 1993). Significant down-cutting of the channel of the Santa Fe River has reduced the potential volume of the alluvial squifix. In many locations, the channel has accured down to the top of the Tesuque Formation. Consequently, stream flow, whether arising from storm-water runoff or upstresm reservoir releases, does not infiltrate but rather flows downstream. This situation pre subsequent recharge of the Tesuque squifer or maintenance of a high riparian water table in the study area.

2.4 Surface Water

This section describes existing physical conditions of the Santa Fe River in the study area. Biological attributes including vagenation, wetlands, fish, and wildtife are discussed in section 2.8. The Santa Fe River watershed encompasses about 160 square mailes (Spiegel and Reldwin, 1963; 190). The headwaten of the river are at Santa Fe Lake, at an elevation of should 11,369 feet near the creat of the Sanger de Cristo Moustnins. The river flows into Cochit Reservoir and the channel continues bodow Cochit Dan to its confluence with the Rio Orande near Cochiti Pueblo.

The portion of the watershed below Nichols Dam The portion of the watershed below Nichols Dam-that drains into the study area comprises about 17,195 acres, or about 17 percent of the entire watershed. Meljor tributaries to the Sants Fe River in the study area include Arrayo de la Juya, Arroyo Tearroo, and Arrayo de Ban Astonia (Figura 1). Sevenul other sensibler survoyo draining the hill slapes on the north side of the river between Carsino Alire and Agua Pris are also tributary to the river in the study area.

2.4.1 Hydrology

Before 1881 there were no narfined water storage facilities on the Sants Fe River (Spiegel and Beldwin, 1963: 172). Divarrious of surface water from the Santa Fe River began with Spanish settlement of the area, which started around 160+ Native Americana inhabiting the area prior Spanish astitement likely did not divert water from the irver for use in growing crops (Spiegel and Baldwin, 1963: 91-92).

Stone Dam was constructed in 1891 on the Senta e River upstream in the cazyon reach of t vor. The reservoir created by the dam filled w

sediment in the span of a few years. Two-Mile Dam was then built at a downstream site in 1894. increasing need for water prompted the construction of Granite Point Dam, about four construction of orange real pain, in 1926. A third structure, Nichols Dam, was constructed between Two-Mile and Granite Point dams in 1943. The crest of Granite Point Dam was raised 1945. The creat of Granus From Dam was raised in 1935 and again in 1947, when the name was changed to McClure Dam. In 1992, Two-Mile Dam was found to be unstable and the dam was subsequently decommissioned in 1994.

Prior to upstream impoundment of stream flow, the Santa Fe River in the study area likely had considerable, sustained flow most of the time. Perennial flow characterized the reach of the river Percennial flow characterized the reach of the river from its source at Santa Fe Lake downstream through downtown Santa Fe (Orant, 2002: 9-10). Surface water flow through the study area reach was likely reliable in most years, as indicated by the historic locations of 11 irrigation diversions on the river along this reach (Spiegel and Baldwin, 1963: 174).

Currently, flow through the study area occurs only in response to storm water runoff or releases from the upstream dams. There are at least 20 discrete storm water drain or arroyo confl Santa Fe River in the study area (Figure 9; Plate 6). These inputs range from small diameter (e.g. 1-foot diameter) pipes, to open arroyo channels, to large storm-water drain culverts.

The estimated 100-year recurrence interval flow in the study area is about \$,000 cubic feet per second (cfr; Lange, 1998). The 100-year recurrence interval flow is defined as a flow with a one percent probability of occurring in any given year. Analysis conducted for the lower segment of the study area estimated the 25-year recurrence interval flow in the Santa Fe River (Le.


the flow with a four percent probability of occurring in any given year) to be 1,800 cfs. Sankfall flow (1.5-year recurrence interval, 67 percent probability of occurring in any given year) was estimated to be 230 cfs (W.J. Miller Engineers, Inc., 2000).

Spiegel and Baldwin (1963: 173) estimated that a flow of 4 of: at the mountain front would infiltrate enjirely into the channel alluvium in the reach downstream to Cieneguita (i.e. current-day Frenchy's Park). It should be noted that this on rate was estimated under conditions of a channel that was not as incised as the current channel and therefore a greater volume of attuvial sediments was available to hold water and allow it to penetrate into permeable sections of the underlying Tesuque Formation.

Nichols Reservoir of 1.5 to 3.0 cfs (three to six arre-feet per day) would be needed to maintain surface flow in the Santa Fe River down to the San Ysidro crossing (M. Hamman, Water Services
Division Director, pers. comm., 10 June 1998). In
spring 2007, a sustained release from Nichols
Reservoir of about 10 cfs maintained flow through the entire study reach (Figure 10). A 10-cfs flo sustained for 24 hours equates to a volume about 20 acre-feet of water.

Senta Fa River Study





Pigare 10. Plow in the Banta Fe River near the NM 599 bridge on 24 May 2007. View is upstream. Photo by J. Pittenger.


2.4.2 Santa Fe River Channel Morphology

Stream channel morphology is controlled by interaction of flow regime, sediment supply, valley slepe, composition of channel bed and benth sediments, and riparian vegetation (Loopeld and Miller, 1956: 28; Leopold et al., 1992: 198; Miller, 1936: 28; Leopold et al., 1992: 198; Leopold, 1994: 5). Valley slope through the study area is about 1.35 percent. Historically, channel materials ranged from pebbly, medium to very course sand in the channel beds to poorly sorted, stightly pebbly, sithy, very fine to medium sand on channel bashs and floodplain (Shrobs et al., 2005). Consequently, the historic channel likely had a meandering plan form with a pool-riffle or gravel bar-smal bed sequence repeated at a spacing of every five to seven channel widths (Leopold, 1994: 28). Such a meandering pattern in a sand-bod stream would be classified as a C5 channet (Rosgen, 1994). A photo of the Santa Pe River near the Alto Street well taken in the ourly 1960 shows an un-incided channel with an obvious meander pattern, which agrees with the presumption of a historic CS channel in the state area (Springerl and Baldwin, 1963: 139). Similarly, a 1969 photo of the sewer line crossing of the river below St. Prancis Drive shows an un-incised channel with an active, fairly wide floodplain (Heggen, 1997: 29).

2.4.2.1 Causes of Channel Incision and Destabilization

A series of busines-induced impacts resulted in dramatic changes in the channel from its probable natural form. First, the bydrograph was radically changed with impoundment of stream flow by upstream reservoirs and loss of the shallow water table from groundwater pumping. These impacts started in the late 1800s and increased in

Plance 9. Locations of storm water inputs to the Santa Fe River in the study area.



Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

magnitude through the late 1950s. Secondly, increased urbanization increased the area of impervious surfaces, which had the double offset of reducing recharge of the shallow squifer and reducing the time of concentration of storm water of rendering frozens to the same and the rendering the time of concentration of storm water runoff flows (Dume and Leopold, 1978: 275-277). Consequently, storm water runoff peak flows increased in magnitude, with an associated decrease in duration. These flows pilics introduced powerful, highly entire bydraulic forces in the channel. Peak storm water flows, coupled with a loss of riparian vegetation that historically made the stream banks more resistant to erosion relative to the stream bed, caused excessive bank session, bed scour and destabilization of the channel.

Third, the City of Santa removed or k existing grade control structures in the river, beginning in 1974, to increase flood-containment acity in the channel (Hoggen, 1997: 9-10). capacity is the channel (Hogger, 1997: 9-10). Grade control structures that were removed included culverts, sills constructed before the 1960s, and rock check dams that were probably built in the 1930s (Hogges, 1997: 9). This campaign of grade control structure removal criticated a period of rupid and dramstic channel degradation. For example, the channel bed below the St. Francis Drive crossing drapped about 12 fast in seven years after removal of downstream conductors of the product of the prod the St. Francis Drive crossing dropped about 12 feet is seven years after removal of downstream grade control structures (Heggen, 1997). This secur effect was reaccrited by sand and gravel mining in the channel, which locally lowered channel base level and initiated upstreamingsting headcutting. Sand and gravel mining in the river channel and floodplain was wridespread from about the CR 62 crossing downstream to the NM 599 crossing. Degradation incised the channel bed through the recent alluvium and into the relatively soft sandstone and conglocenstre at the top of the Teasuage Formation in many the top of the Tesuque Formation in many locations through the study area.

Santa Fe River Study

These impacts imposed new conditions to which the channel must adjust. Channel incision has been defined us a morphological expression of an imbalanced condition where sediment transport capacity exceeding sediment supply. To re-establish dynamic equilibrium, the channel undergous changes in along and cross-rectional area (Harvey and Watson, 1986). An characteristic sequence of changes in channel cross rection occur following bed degradation (Harvey and Watson, 1986). The major stages in channel vicinity of the channel with the channel with channel strength of the control of the changes in channel strength of the channel s

As channel widening slows due to increased cross-sectional area and reduced sediment transport capacity, a Type IV channel form develops. This channel form is characterized by ncreased sediment deposition on berms or necent floodplains, decreasing channel slope, and increasing width-to-depth ratio. Finally, a state of increasing width-to-depth ratio. Finally, a state of dynamic equilibrium develops and Type V channel form develops. Type V channel is characterized by a relatively high width-to-depth artic, flattening of channel slope, and continued aggradation and development of a floodplain. This channel type is in dynamic equilibrium, where erasion and deposition processes are in https://dx.org/10.0045/97. balance (Briggs, 1996; \$2).

2.4.2.2 Classification of Stream Segments in the Study Area

Mentification of the evolutionary stage of channel reaches in the study area was conducted by applying the Rosgen stream assessment technique and morphological classification system (Rosgen, 1996). The Rosgen stream morphological classification system lend with the reachest classification system lends itself well to assessing channel evolution status, as it provides standard method to compare stream segments to the probable natural, equilibrium form (i.e. the Type V channel discussed above).

Level II stream assessments were conducted by J. Pittenger of Blue Earth Beological Consultants, Inc., in 1999 at two locations in the study area frenchy's Park and the State Trust land upstream from NM 599) and one location immediately upstream from the study area (the reach between St. Francis Drive and Camino Alire). These ats involved surveying the slope of the assessments involved surveying the slope of the channel, determination and measurement of bankfull channel width, measurement of stream substrate particle size distribution, and measurement of other channel morphology features (Rosgen, 1996: 5-15 through 5-29). Stream types delineated during these assessments

- F4b stream type in the reach between St. Francis Drive and Camino Alire;
 P5 stream type at Frenchy's Park;
 B5c stream type in the reach from NM 599 upstream to the meanders below the old Leeder river crossing;
 C5 streamtype in the meandering reach below the old Leeder river crossing;
 C5 stream type above the old Leeder river crossing;

These delineations were made prior to the restoration work at the State Trust reach, which restonation work at the scatte frum reach, water resulted in removal of the Leeder stream crossing, and the project from St. Francis Drive to Camino Alire. The 1999 delineation data provided a basis or key for the following broad level I assessment constructed in the study area (Rosgen, 1996: 4-20

The following level i assessment was conducted by walking the entire study area and recording observations of plan-view morphology (i.e. extent of meandering), channel width and depth at bankfull stage, channel slope, and bod feature. These field observations, along with the 1999 delineation data, were then used to interpret and delineate every account on a garial photography.

The following delineation is presented from the The religioning a cimication is presented from the upstream end of the study area, starting at Camino Alire bridge, and proceeding downstream to the NM 599 bridge. The thalwag of the river channel in the study area was first mapped and subdivided into 200-foot intervals to facilitate location and description of stream segments. The 200-foot interval stations start at the NM 599 bridge (station 0+00) and proceed upstres Camino Alire bridge (station 378+80).

Three stream types were delineated in the study area: B, C, and F. The C and F type channels were further subdivided according to dominant channel materials; either gravel or sund (Table 1). The B stream type in the study area had gravel-dominated bed material. Over half of the study dominated bed material. Over half of the study area (34 percent) was characterized by C-type channel, which was defineated in five reaches. The F-type channel composed 38 percent of the river in the study area and was defineated in six reaches. One reach was defineated as B-type channel, which composed five percent of the river in the study area (Table 1; Plate 6).

Santa En Blose Shelv

Page 18

U.S. Army Corps of Engineers, Albuquerque District

Figure 11. Delineation of channel segments in the study area using the Rusgen classification system




Table 1. Rosgen stream type classification of channel segments in the study area.

			Start	End
SFR-1	B4	1,767	360+3 5	376+02
SFR-2	F4	7,123	289+12	360+35
SFR-3	C4	4,614	242+87	289+12
SFR-4	F4	633	234+64	242+97
8FR-5	C4	1,000	224+64	234+84
SFR-6	F4	584	219+00	224+64
SFR-7	Ø	1,458	204+42	219+00
SFR-8	F4	1,509	159+33	204+42
SFR-9	C5	2,684	162+89	189+33
SFR-10	F5	2,542	137+27	162+69
SFR-11	F6	1,799	119+28	127+27
8FR-12	C5	11,928	0+00	119+28

The B stream type is moderately entrenched with moderate bandfull width-to-depth ratios and a relatively steep gradiens (2 to 4 percent). The B-type channel in the study area has a bed dominated by garvel with some send and cobbles. In the study area, the 84 stream segment is bounded on the downstream end by a grade courted structure. This stream type was delineated in segment SFR-1, which is a 1,767-6oot reach downstream from Carnino After (Figure 11). Gabbino banker walls line portions of the benks in this meach. Decreatation is not occurring in the B-title meach. Gabien beaket walls time portross of the beats in this reach. Degradation is not occurring in the B-type channel segment in the study area. Furthermore, rapid channel widening is not occurring, as evidenced by vegetated banks and absence of rew, vertical cause (Figure 12). Therefore, following the preceding discussion in section 2.4.2.1, the probable channel evolution stage of segment SFR-1 is IV or V, which is nearing equilibrium condition.

The C stream type is less entreached than the B stream type, has a relatively high bandfull widthto depth retio, and has moderate sinussity. This
stream type was delineated in five locations in
study area: segments SPR-3, SPR-5, SPR-7, SPR-9, and SFR-12 (Figure 11). Begment SPR-3 is
notated in the vicinity of SHR Road and consists
of a 4.614-foot reach with a relatively welldeveloped floodplain (Figure 13). The
downstream end of segment SPR-5 is marked by
the Veroda de San Antonio road crossing, which
compared to many other reaches in the study area
(Figure 14). Segment SPR-7 is a borderlins C4F4 chemnel and extends from the Do Antonios
crossing downstream to the San Ysiddo crossing. cossing downstream to the San Yaidro crossing

Channel bed composition shifts from gravel-dominated to sand-dominated in segment SFR-9, which is the San Yaidro River Park restoration which is the Saa Yaidro River Park restoration reach from the large grade control structure downstream to CR 62 (Pigare 11). Finally, segment SFR-12 is a long reach extracting from near the Cottonwood Mobile Home Park downstream through State Trust land to the NM 539 bridge. The lower portion of this segment in particular has a relatively wide floodplain and moderate simuosity (Pigare 15).

The C-type channel in the study area ranges in evolutionary stage from reactice that are still actively widening (Type III) to those that are nearing equilibrium (Type IV; Harvey and Watson, 1986).

Santa Fe River Study

Page 17

U.S. Army Corps of Engineers, Albuquerque District



Figure 12. Stream type B4 below Centha Alire bridge. Arrow indicates field determination of beakfull stage. View is upstream, with the bridge in the beckground. Photo by J. Frittenger, 28 May 2007.



Figure 13. Stream type C4 near the Siler Road dead-end. Relatively well developed floodplain is visible in center-left. View is upstream. Photo by J. Pittenger, 24 May 2007.

Type III (i.e. disequalibrium condition) C-channel reaches are characterized by rapidly eroding, often steep-walled bases. Examples of these are found in segment SFR-12 upstream from the Cottonwood Mobile Home Park and in segment



Figure 14. Stream type C4 above the Verada de San Autonio crossing. Note the wide, accessible, undeveloped floodphain area on the right and the absence of channel entrenchment. View is downstream. Patot by J. Pittenger, 26 June 2007.



Figure 15. Stream type CS at the lower end of the study area. A relatively well developed floodplain, moderate simuosity, and absence of entreachment characterize the channel in this reach. View is upstream. Photo by J. Pittenger, 24 May 2007.

Senta Fa River Study

ITHE P stream type is characterized by an entrenched channel, moderate sinuosity, and a high bankfull width-to-depth ratio. Six stream segments were delineated in the study area as P channel, two of which are configuous (Table 1). Segment SPR-2 is the longest reach of F-type channel is the study area (Table 1; Figure 11). This segment extended from the arroye confluence near Nix Stables upstream through Frenchy's Parkto station 360-13 (Figure 17). The downstream end is marked by a grade control structure (GCS-7 in Plate 6). Three relatively abort segments of F4 channel were defineated from new Siler Road downstream to below the San Ysideo crossing (Table 1; Figure 11). Finally, two contiguous segments of F5 channel were delineated from the CR 62 crossing downstream to near Cottonwood Mobile Home Park (Figure 11).

The F-type channel segments have the greatest degree of disequilibrium condition in the study area. Some of these reaches, such as segment SFR-11, appear to be closer to equilibrium as indicated by vegetation establishing slong the toe of croded terrace escarpments. The F-channel character of segment SFR-4 may be a recent development resulting from damping of fill and marked constriction of the floodplain (Figure 18). Other reaches, such the upper pert of segment SFR-10 (below CR-62) and SFR-2 (upper reach in the study area), are in early stage of adjustment and are rapidly increasing channel width as indicated by crotion of high, vertical banks (Figure 19). The formation of a few lateral bars in these pertions of SFR-10 are indicative of some degree of aggradation, which implies that bed degradation has slowed in this segment. These field indicators suggest that these reaches are at the Type II or III channel evolution stage (Harvey and Watson, 1936). Under current confliction of grade control, both ranches will likely continue to this stage that the study of the grade control structures influences these reaches greatly and if they are compromised the channels would begin degrading.

U.S. Anny Corps of Engineers, Albuquerque District




Figure 16. Type III (disequilibrium) C5 channel. Reach is located in segment SFR-7. Arrows indicate actively eroding bank and zone of rapid channel widening. View is downstream toward the San Ysidro crossing. Photo by J. Pittenger, 26 June 2007.



Figure 17. Stream type F4 upstream from Frenchy's Park. Coarse bed material and psucity of aggradation indicate the bed in this reach is still degrading.
View is upstream. Photo
by J. Pittenger, 15 June
2007.

U.S. Army Corps of Engineers, Albuquerque District



Figure 18. Channel-narrowing fill along right bank in segment SFR-4. View is downstream. Photo by J. Pittenger, 26 June 2007.



Figure 19. Deeply-entrenched P5 channel entrenched P5 castanes below CR 62. View is downstream. Photo by J. Pittenger, 19 September

2.4.2.3 Grade Control Structures in the Study Area

Seventeen grade control structures are located in the study area (Table 2). Grade control structures are concentrated in the upstream half of the study area (Figure 20; Plate 6).

Table 2. Grade control structures in the study area. Types are GAB = concrete-capped gabion wall, ROCK = rock wall, CW = concrete wall, CSS = concrete step structure, and LWC = concrete low-water crossing.

			PLESS RESELVED
GCS-1	GAB	1	99+87
GCS-2	CSS	15	182+69
GCS-3	CW	2	189+34
GCS-4	CSS	12	204+87
GC8-5	LWC	6	224+68
GCS-8	GAĐ	4	284+00
GC8-7	GAB	3	289+11
GCS-6	GAB	2	294+34
GC8-9	GAB	3	298+35
GC8-10	GAB	1	302+00
GC8-11	GAB	2	309+07
GCS-12	ROCK	4	344+00
GC8-13	GAB	2	360+35
GC8-14	GAB	3	368+48
GCS-15	GAB	3	387+61
GC8-16	GAB	2	369+85
GC8-17	GAB	3	377+29

Two of the grade control structures provide major control, in that their effect pernists for considerable distances upstream. The CR 62 crossing (GCS-2) has a downstream drop of about 15 feet while the San Yaidro crossing (GCS-4) has a downstream drop of both 12 feet. The Vereda de Sen Antonio low-water crossing (GCS-5) is also a significant control of bed elevation in the

2.4.3 Water Quality

Surface water flow in the study area occurs in Surface water flow in the study area occurs in response to upwercam reservoir releases and storm-water runoff. There are no water quality standards established for the Sants Fe River in the study area. When there are surface water flows through the study area, water quality is affected by high levels of asspended sediment, chemical contaminants from urban sterm-water runoff, sad-trash and debrie that is dumped into the river throughout the study area (Orant, 2002: 26). The throughout the study area (orant, 2002: 29). Itself or irver is the study area was assessed in 2004 and found to not support aquatic life, secondary contact, and wildlife habitot designated used to polychlorinated biphenyls (PCB) in the water column during forom water tunoff events (New Munico Environment Department, 2007: 225).

Urban storm-water runoff is typically contaminated with sediments, nutrients, microbes, toxic metals, and organic compounds (Makepeace et al., 1995; Pitt et al., 1995). Runoff from the et al., 1995; Fit et al., 1995). Runoff from the rond network in the study were actehnent likely in a major source of pollutants in storm water. Pavement surface wear, brake lining wear, tire wear, fuel and oxhaust, oil, grease, hydraulic fluids, and engine and parts wear generate pollutants unch as sulphates, subsettos, copper, nickel, chromium, particulates, rubber, zinc, lead, PCB, and petroleum compounds (Forman et al., 2003: 202-206).

Santa Fa River Study

Page 24





Figure 21. Grade control structure GCS-2 at the CR 62 crossing. View is upstream. Photo by J. Pittenger, 24 May 2007.





Figure 22. Grade constructure GCS-5 at the Vereds de San Antonio crossing. View is upstream. Photo by J. Pittenger, 26 June 2007.



Figure 20. Grade control structures in the study area.



Santa Fe River Study

Page 25

U.S. Army Corps of Engineers, Albuquerque District

2.5 Soils

Six soil mapping units occar in the study area (Table 3; Natural Resources Conservation Service, 2007). However, only two of these soil mapping units are common. These are Riverwash and Riovists gravelly loamy sand, which together comprise over 96 percent of the soils in the study area. The remaining four soil mapping saint each comprise less than two percent of the study area and are evoded terrace remainst in the area of river influence (Figures 23-34).

Table 3. Soil mapping units in the study area.

Soil Mapping Unit	Acres
101 Zozobra-Jaconita complex	0.12
106 Pits	0.49
107 Rhysrwath	64.30
112 Riovista gravelly loamy and	16.76
113 Delvalle-Liran land complex	1.31
116 Argente-Urban land-Orthenta complex	0.96
Total	82.67

The Rivacwash mapping unit consists of gravelly course sand and gravelly sandy loans formed from alluvium. These soils are frequently flooded and cocur in the river channel and its nascoss floodplain in the study area. The soils are excessively drained and have moderately high to high capacity to treasmit water. Scharated transmissivity of Rivarwash soils ranges from 0.37 inches/hour to 1.98 inches/hour. Consequently, available water capacity is very low (cc. three inches).

Riovista gravelly sandy loans occurs on floodplam and valley floor areas and is formed from alluvium derived from granite, gnates, and echiat. These soils are also accessively dutined and have very high saturated hydraulic transmissivity (6.0 to 20 inches/hour). A veitable water holding capacity in Riovista sandy gravelly loam is about 1.7 inches.

The remaining floar minor soil mapping units countie of soils formed from allavium that occur on croded fan remeants or stream termoer. These soils are gravedly occurs under Joanne to sandy loams. These soils are well drained but have slightly higher water holding capacity compared to the Riverwath and Riavium gravelly learny sand soils because of a generally higher percentage of loam.






Figure 23. Soils in the study area, station 0+00 to 32+00.



Figure 24. Soils in the study area, station 32+00 to 64+00.

Page 28



Figure 25. Soils in the study area, station 64+00 to 32+00.



Figure 26. Soils in the study area, station 102+00 to 138+00.

Sente Fe River Study

Page 29

U.S. Anny Corps of Engineers, Albuquerque District



Figure 28. Soils in the study area, station 176+00 to 196+00.

Figure 27. Soils in the study area, station 138+00 to 176+00.



U.S. Army Corps of Engineers, Albuquerque Di




Figure 38. Soils in the study area, station 228+00 to 264+00.

Figure 29. Soils in the study area, station 196+00 to 228+00.





Figure 31. Soils in the study area, station 264+00 to 298+00.



Figure 32. Soils in the study area, station 298+00 to 334+00.

Sente Fe River Study

Page 32



Figure 33. Soils in the




Figure 34. Soils in the 364+00 to 378+02.

Santa Fe River Study

Page 33

U.S. Army Corps of Engineers, Albuquerque District

2.6 Air Quality and Noise

2.6.1 Air Quality

The Clean Air Act of 1970, as amended, established National Ambient Air Quality Standards (NAAQS) for six criteris air pollutants: come, airbonne particulates, carbon moneciach airrogen dioxide, sulfur dioxide, and lead. If the concentrations of any of these six pollutants exceeds the standard, the area where the exceeds are standard the area where the record and the standard the area where the record and the standard the area where the cuterions of the standard the st exceedance occurs is considered non-striament for that pollutant. No violations of the NAAQS have occurred in Santa Fe County, which is classified as attainment for all six criteria air pollutants by the Baviroumental Protection Agency (U.S. Environmental Protection Agency, 2007).

2.6.2 Noise

In considering potential effects of increased noise in considering potential effects of increased noise levels, sensitive noise receptors are identified in a study wes. Sensitive receptors include, but are not limited to, homes, lodging facilities, hospitals, parks, and undeveloped subards areas. Sensitive receptors in the study area homes, parks, and

Existing noise levels throughout the study corridor are both natural and manmade, including rushing water (when the river is running), rustling leaves, domestic and wild animats (e.g. birds, dogs, insects), human voices (e.g. children at play, trail user talking on a cell phone) and activities (e.g. hammering, unloading building materials), and whiches and other machines (e.g. bisycles crossing metal bridge, lawn mowers, power saws, erawd-smining conjourners.

In the eastern-most portion of the study are so a a summer weekday merning, there was surprisingly little noise filtering into the study area from surrounding streets and homes along some portions of the corridor. In the central portion of the corridor, sounds from businessee can be heard, such as fines at a door measurfacturing plant and tracks being founded, as well as human voices and tractic on Agus Pris Street and side streets. At the traffic on Agus Fris Street and side streets. At me west end of the corridor, the most prominent sound is traffic on N.M. Highway 599.


2.7 Ecological Setting

2.7.1 Plant Communities

Plant communities in the study area were mapped using the community-structure (C-S) classification scheme developed by Hink and Ohmart (1984). This classification combines identification of community dominants in the tree and shrub strats with the structural character of the stand being delineated, where structural character is defined as the variation in foliage density with height above the ground surface.

Six structure types are used in the classification. These range from structure type I, characterized by an overstory canopy provided by maters trees (i.e. 50 to 60 feet tall) and understory foliage to (i.e. 50 to 60 feet tall) and understory foliage to type VI, characterized by apares herbaceous and shrubby vegetation (Figure 35). A seventh structure type X, was added to the classification to describe lacking woody dominants and with foliage restricted to three feet above the ground and lower. U.S. Army Corps of Engineers, Albuquerque District

Figure 35. Vegetation structure types in the Hink and Ohmart classification scheme. The figures are graphical representations of the variation in foliage density with bright above the ground surface. Excerpted from Figure 3 in Hink and Ohmart (1944: 38).



Plant community types were classified using a combination of 13 codes for decrinent or co-dominant species or cover type. Six of the codes were for native woody or suffluencent plant species: C for cottonwood (including Rio Grands, earrowleaf, and Inco-leaf). CW for copote willow, OW for Coodding's willow, I for one-seed jumper, LO for New Mexica locust, and RB for tubber rabbitrouts (Table 4). Another five codes were for non-native woody plants: HL for homey-

locust, RO for Russian olive, SC for saltcedar, SE for Siberian chm, and TH for tree-of-heaven (Table 4).

Two miscellamous cover classes were used for area lacking woody vegetation. The code BARE was used for area with sparse berbacous cover and greater than 70 percent burs ground. The code HBRB was used for areas dominated by

Dominant herbaceous species in areas delienated as HERB included hairy golden-aster (Heterorheca willien), horseweed (Conyea canadanati), asmooth acryo (Heterorheca willien), horseweed (Conyea canadanati), asmooth acryo (Heterorheca willien), trough cockdebur (Kanshisen strumarinen), prickly lettace (Lachaca aeriole), burra rayweed (Andropsia acconsilicarpsi), Russian-shietis (Saleola Iraque), white sweet-clover (Meliliona albus), sorred with white sweet-clover (Meliliona albus), sorred deliberation (Chunathy and Chunathy (Eriognams polycladens), caladigue (Russes (Menoraspalus), Chardinas (Meronas Incirorum), Carolina lavegrass (Eragrossis pectinacae vas peccinacae), and foxtail bartey (Hordenss Jabonans).

Major factors influencing the ourrent condition of riparian vegntation in the study area are 1) significantly reduced surface water flow; 2) has of the shallow athwais aquific; 3) massive bed degradation throughout the reach initiated in the mid-1976 by removal or lowering of grade control structure; and 4) securing peak flows associated with stem-water ranceff that are new contained within a narrow, entreached valley throughout most of the study area. Because of these factors, riparian vegetation is specially others throughout the study area and where it is found it to typically characterized by early successional stages.

cont	SPECIESTONES				
BARE	Mostly bare ground with scattered herbaceous plants				
HERB	Harbaceous vegetation				
С	Rio Grande, narrowleef, andier lance-leef cettomeood; lance-leef is a hybrid between Rio Grande and narrowleef (Populus distributes wistizanti, P. angueticitis, andier P. x acuminate)				
CW	coyote willow (Saffx exigua)				
GW	Goodding's willow (Saffx gooddingff)				
J	one-seed juniper (Junipenz monospame)				
LO	New Mexico locust (Robinie neomexicena)				
RB	rubber rabbitbrush (Ericamerie neuesces)				
HL.	honey-locust (Ginditale triacanthos)				
RO*	Russian otive (Geoegnus angustitolis)				
SC*	sakouder (Terrech chinansis)				
se•	Siberian eim (Ukrus pumile)				
TH	tres-of-heaven (Allantitus attissima)				

U.S. Army Corps of Engineers, Albuquerque District

Table 5. Coverage, in acree, of plant community-structure (C-S) types mapped in the study area. A slash "/" separates overstory dominants (left of slash) from understory woody dominants. Codes separated by communication in strain. For example, C-S type C/CW, RB is a plant community with an everstory dominated by cottonwood and an understory shrub strate dominated by coyete willow and rubber rubbilihored.

	1251775						orale at a series
		N		٧١	K	EXAMINATION OF THE	
SARE					37.00	27 M	45.76%
HERD		***	-		444	4.80	8.14
<u>e</u>	0.92	J.36	9.27	8.10		0.63	1,61%
c.ew			4.05		1	8.40	811%
C.OW/CW			8.04		-	1.14	1.11%
0.01		8.27				0.27	8.53%
DIRECW		131	3.01	D.M	-	1.48	1.70%
CICM		1.01	0.41	1.00		3.42	6.13%
CICW.84				1.34		1.04	1 25%
C/RD				2.13		2.53	3,86%
CW				1.86		2.91	3.65%
CW.16/R1				1.69		0.04	6.75%
HL				1.05	-	48.0	0.05%
HLSE/CW				9.86		0.00	8 47 %
HLTH.BE			0.11	-		0.11	4.15%
JMS				1 12	-	5.12	1 16%
10			-	8 14		8.34	0.17%
89				10.03		18 23	21 73 W
ac.Rp.cW/RE	 -			0.21		1.25	1 20%
10		9.76	0.01	1.16		1.03	2,33%
10.0		0.10		6.00	-	0.10	D 28%
SE RD.CW/RD				0.12		8.32	0.30%
RB/CW		9.41			 -	0.41	0.51%
BEICH			+	3.42		9 43	4.14%
		-	0.02			0.01	0.02%
TH,86		+=	1.24			8.21	9.29 %
TH.GEMD		4.12	1112	34,83	42.50	92.67	
	1.02	1.12	1,55%	41.12%	51.32%	1	

Over half (51.32 percent) of the shady area consisted of bare ground areas or herbaceous plant cover (C-S types BARE X or HERB X; Table 5). Structure type VI stands, which are plant communities with a majority of the foliage least han 10 feet high, made up another 41 percent of the study area [Table 5]. Structure type VI stands were highly variable in terms of dominants. A total of 17 differents community types with this structural stage were delineated in the project area framework. Table 5] Structure type II repartition, which is characterized by mature trees (cf. Figure 35) comprises only 0.03 percent of the study area and consists of a single C II stand along the left bank between CR 62 and Cottonwood Village.

Aside from the single C II stand described above, plant C-S types dominated by native woody apocies included: C,GW VI; C,GW/CW VI; C/CW VI; C/CW VI; C/CW VI; C/CW B VI; C/CR B VI; C/R B VI; and CW VI (Table S). These native riparian C-S types are concentrated in three main areas:

- the lower end of the project area upstream from NM 599 to just upstream from Cottonwood Drive, with most occurring in the section from the old Leader backs plant lease downstream to the NM599 bridge (Plate 1);
- the Siler Road area, from the Vereda de San. Autonio crossing (Plate 3) upstressu to the lower end of the Boytan property (Plate 4); and
- a small area near the confluence of Arroyo Torroon downstream from the Camino Alire bridge (Plate 5).

These patches of native woody riparian ranged in structural type from small trees (type IV and V; Figures 36 and 37) to stands of saplings or willow shrubs (type VI; Figure 38). Vegetation patches dominated by native woody rigarian species totaled 11.67 scres, or about 14 percent of the vegetation in the study area.

Sites of native woody riparian vegetation ortablishment are characterized by stream reaches with sediment deposition on lateral or point berr adjacent to the active channel. The Siter Road area and the downstream portion of the study area both are relatively long stream reaches that appear to be at or near equilibrium state, with developing meander patterns and floodplains. The absence of substantial establishment of sative woody riparian vegetation in most other portions of the study area are indicative of disequilibrium channel conditions.

Rubber rabbitbrush vegetation on floodplain and ber sites is common in the study eren (Pigure 38). This vegetation, unspeed as C-S type RB VI, maken up about 22 percent of the vegetation in the study area (Table 5).

The most common non-ustive woody plant in the study eras is Biberian olm. Vegetation stends with Siborian elem totaled about nine sares, or 11 percent of the vegetation is the study area. Other non-native woody species were more localized in their distribution in the retary area and included femory-locuse, tree-of-knewes, and salteedar (Figure 39). Salteedar was found as a dominant is not you location, between stations 64-00 and 74-53, below the Customwood Drive crossing and 74-53, below the Customwood Drive crossing at this location in spense napling stands with Russian olive and coyote willow (Plate 1).

Senta Fe River Study


Page

U.S. Army Corps of Engineers, Albuquerque District



Figure 37. A C/CW VI stand on the State Trust reach above NM 599, 28 May 2007. Vegetation restored in this reach is initiating channel sarrowing. Photo by I.

Figure 36. A C IV stand near the Siler Road doad end, 26 June 2007. Trees are lance-leaf cottonwood Photo by J. Pittonger.






Figure 38. Rabbitbrush (RB VI) vegetation on a floodplain site downstream from Siler Road, 24 May 2007. View is downstro Photo by J. Pittenger



Figure 39. Stand of Siberian elm, honsy-locust, and tree-of-heaven below Camino Carlos Rael, 26 June 2007. View is stream. Photo by J. Pittenger.

Sente Fe River Study

Page 40

U.S. Army Corps of Engineers, Albuquerque District

2.7.3 Fish

The historic occurrence of fish in the Santa Pe The historic occurrence of fish in the Santa Fe River through the study area is poorty documented. The most notable record is of American sel (Anguilla restruta), an anadromena species, which was collected from the Santa Fe River somewhere between present-day Preuchy-Field and Augu Friso or J Lone 1931 by the notable field and Augu Friso or J Lone 1931 by the notable high selection of the Santa Fe River are accessioned at the Smithnoisus Museum (USNM 16734). No other fish collection records are available for the study area. Trout (Oncorhynchus sp.) historically area. Trout (Oncorhynchus sp.) historically fish collection records are available for the study area. Trout (Oncarhynchus sp.) historically occurred in the Sants Fe River in the downtown area (Grant, 2002: 10) and may have been at least a seasonal part of the ichthyofaums of the study area before major changes in the hydrograph and habitat of the river. Rio Grande sucker (Pantostean plebetiss) occurs downstream from the study area, where perensials flow in maintained by wastewater treatment plant effluent (I. Pittengen, unpubl. data). Rio Grande sucker (Illuent (I. Pittengen, unpubl. data). Rio Grande sucker (Illuent (I. Pittengen, unpubl. data). Rio Grande sucker (Illuent (I. Pittengen, unpubl. data). Rio Grande suckers (Illuent (I. Pittengen, unpubl. data). Rio Grande suckers (Illuent (Illu Pittenger, unpubl data). Rio Grande sucker and fathead minanew (Pinephales prometal) were collected from the river further downstream, in the vicinity of La Cienega, in 1958 (Kanasa University museum records KU 4260 and KU 4262, respectively). Other species that potentially may have occurred in the Santa Pe River in the study area include longaone dace (Rhistichtyse cataractea), and Rio Grande de tub (Gila panelo). These species are common in north-central New Maries present the same descriptor une and the same descriptor une an Mexico streams at the same elevation range as the Santa Fe River in the study area (af. App odia).

2.7.4 Wildlife

The study area provides limited habitat for tearestrial wildlife, due to the narrowaces of the corridor and paucity of vegetation. Thirty-one bird species were obsaved during the field surveys conducted from May through September

2007 (Table 6). Bird species richness was highest in habitats with structural diverse vegetation, particularly the reach from Camina Alire downstream to be old stream gage, the vicinity of Siler Road, and the downstream end of the study area between NM 599 and Cottnawood Drive Scaled quail were relatively common in undoveloped portions of the study area and around horse stables adjacent to the river contidor.

Mammala observed in the study area during the field savery included desert cuttonial (5)-vidagus auduborti), black-tailed jackrabits (Lepus cutifornicus), Camaisson partin dog (Cymonys garninos), striped stunk (Maphitis maphitis), and feral est (Felir catus). Prairis lizard (Sectioperus undustaus) and plateau striped whipsail (Camaidophorus velox) were common reptiles in the study area.

Animal species that may potentially occur in riparian and aquatic habitats in signilar landscape actings as the study area in Santa Fe County are listed in the Appendix.

2.8 Endangered and Protected Species

No federal or state listed species were observed in the study area. Suitable habitat for the federal endangered Southwestern Willow Plycatcher (Empideness traitfill extiluous) is not found in the study area. Suitable habitest is also lacking in the study area. Get Beld Bagle (Hallacetus lemococphahus).

Page 42

2.7.2 Wetlands and Water of the U.S.

No wetlands are mapped in the area and none were documented during the field survey conducted in 2007. However, wetlands historically occurred along the river in the study area, most notably in the vicinity of Frenchys Park (historic sits of Ciceneguits Spring) and Agua Fria (Spieget and Baldwia, 1963: 132) and st esveral pueblo sites including Lor Patacies, Pino, and Pueblo Quemado (Spiegel and Baldwia, 1963: 1963: 1963).

Remnant indicators of hydric solls were found in a vertical cut-bank below CR 62 at Agua Fria during field surveys in 2007. These indicators included a thin organic carbon strats overtaid by a band of iron oxide staining (Figure WET). These features were present in the cut bank about 10 feet above the ourrour channel bad.

The Santa Fe River is considered a water of the United States. Consequently, drodge and fill activities conducted below the ordinary high water mark in the river are regulated by the U.S. Army Corps of Engineers pursuant to section 404 of the federal Clean Water Act and section 10 of the Rivers and Harbors Act.




Figure 48. Resmant by drie soil indicators in a cut bank below CR 25 at Agua Pris. Inset photo "A" shows the location of the indicator status about 10 feet above the current channel bed. Photo "B" shows the thin black carbon strate overfaid by iron oxide-stained sediments. Photo by J. Pittenger, 19 September by J. Pittenger, 19 September 19. Pittenger, 19 Sept

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

Table 6. Bird species observed in the study area during senteer 2007 field surveys.

	Charactus voolena
Scaled Quell	Cultivapia equamate
Turkey Vulture	Catherina sure
American Kestrel	Falco apervedut
Rock Dove	Columba fivie
White-winged Dave	Zenaldo asiatica
Mouning Dove	Zanaida macroura
Black-chinned Hummingbird	Architectus alexandri
Cassin's (Cosbird	Tyrennue vocFerens
Black Phoebe	Sayomia nigricana
Say's Phoebe	Seyomia saya
Northern Rough-straged Swallow	Stelgidopteryx serripannia
Rank Suraktur	Ricaria riperia
Barn Swalkow	Himmito ruellos
Ciff Swalow	Petrochelidon pyrrikonota
American Crow	Convus brachyrhymchos
Common Raven	Convue corex
Servick's Wren	Thryomenee bewickli
American Robin	Turtius migralorius
Northern Mockingbird	Mirrus polyalottas
Curve-billed Threater	Toxostome currirostre
European Starling	Stumus vulgaris
Yellow Warbler	Dendroica patechia
Stack-throsted Searrow	Amphicpiza bilineata
Bullock's Orloin	faterus bullocidi
Western Meadowark	Sturnelle neglecte
Brown-headed Combird	Mobthrus ater
Canyon Towner	Pieto fuscus
Scotted Towner	Pipilo maculatus
Black-handed Groebeak	Pheuticus melanocaphalus
House Finch	Cerpodecus mexicanus

Sante Fe River Study

2.9 Socioeconomic Environment

The Santa Fe River study area is located in Santa The Santa Fe River study area is located in Santa Fe County, New Mexico. Approximately 1.5 miles of the 7.16-mile study area are within the city limits of Santa Fe, New Mexico. Santa Fe with a January 2007 population estimate of 68,359 (City of Santa Fe, 2007), is the capital of New Mexico and the test of the tatte povernment. It is a full-arcvice community with emergency services (fire, police, medical), a bospital, public and crivites exhoust churchs museums therefore the control of the country of the santa for the country of the country and private schools, churches, museums, retail and private schools, churches, museums, retail stops, restaurants, and other services. According to Santa Fe Trends (City of Santa Fe, 2007), the city's largest employment sectors are government (29 percent), retail/wholesale (15 percent), hospitality/food service (13 percent), health cars/accial assistance (10 percent), and construction (7 percent). The remainder of the study area is located in unincorporated Santa Pe County. As of July 2006, the County's estimated population was 142,407 (U.S. Census Buresu, 2007e). The study area includes portion of Santa Fe County Census Tenests 8, 12.01, 12.02, and 12.03. (U.S. Census Buresu, 2007e). Using Census 2000 data, some population demographics of these four cessus tracts are compared to those of the City of Santa Fe, Santa Fe County, and the State of New Mexico in Table 7. The data show that the combined population of the first Santa Fe County census tracts that include the study area have a slightly lower percentage of white pearons, and conversely, fewer ethnic minosities, than are found in the City of Santa Fe, Santa Fe County and the State of New Mexico. These four census tracts also have a much higher percentage of percons (74 percent) identifying demarked as allispassic or Latino than do the total populations City of Santa Fe, Santa Fe County, or the State of New Mexico, all three of which have less than half of their populations identifying themselves as Hispanic or Latino (Table 8).

Table 7. Comparison of racial demographics for the State of New Mexico, Santa Fe County, City of Santa Fe, and combined data for Santa Fe County Census Tracts 8, 12.01, 12.02, and 12.03, which include the study area. Data are from Census 2000.

				in Fo Count Cines Tracts
Total population	1,819,046	129,292	62,203	19,10
White	66.8 %	73.5 %	76.3%	B2.4 1
American Indian	9.5 %	3.1 %	2.2%	2.2 1
Back or African American	1.9 %	0.6 %	0.7%	0.8
Asian	1.1 %	0.9 %	1.3%	0.4 5
Native Hawsing/Pacific Islander	0.1%	0.1 %	0.1%	0.2
Some other race	17.0 %	17.7 %	15.3%	29.7
Two or more reces	3.6 %	4.1 %	4.2%	4.6

U.S. Consus Bursau (2007a)

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

2.10 Land Use, Recreation, and Aesthetics

Land use, recreation, and seathetics in the study area are first summarized in this section and then followed by more detailed descriptions organized by river reach. For the purpose of this discussion,

- 1. Camino Alire to Camino Carlos Rael (ca. 1.5
- miles);
 2. Camino Carlos Rael to CR 62 (ca.2.6 miles);
 3. CR 62 to Cottonwood Drive (ca. 1.5 miles);
- 4. Cottonwood Drive to NM 599 (ca. 1.5 miles).

The 7.16-mile Sants Fe River study area, defined as the area ecologically influenced by theriver (cf. section 1.1), includes private, federal, accountly, and city lands comprising about 83 acros (Tables 10). These various ownerships are depicted in Figures 41 through 45. Approximately 1.48 miles of the river study area within the Sants Fe (city limits, and the remainder (f.68 miles) is within unincorporated Sants Pe Country.

As seen in Figures 41 through 45, in general, lands adjacent to the study area that are under county jurisdiction are less-densely developed than lands in the city segment. The city-county boundary is approximately at the Camino Carlos Rael crossing. This is particularly true the farther one moves downstream (away from the city) and on the north side of the river in the unincorporated county reach. county reach.

Table 10. Acresge by land ownership in the study area.

City of Santa Fe	0.44	0.53
Santu Fe County	33.12	39.92
Sinte of New Maxico	3.19	3.04
Bureau of Land Management	1.17	1.41
"Common Area"	8.33	10.04
erge of the state of	and the second	
Private	38.72	44.26
### 47 FOR SALE		

U.S. Army Corps of Engineers, Albuquerque District

Table 8. Proportion of Hispanic or Latino and non-Hispanic or non-Latino residents of the State of New Mexico, Santa Fe County, City of Santa Fe, and combined data for Santa Fe County Cansus Tracts 8, 12.01, 12.02, and 12.03, which include the study area. Data are from Census 2000.

an aftern	New Marks	Carry 1-	//盘/类	Santa Fo December Common Tracto
Total population	1,819,046	129,292	62,203	19,104
Hispanic or Latino	42.1%	48.0%	47.8%	74.3%
Not Hispanic or Latine	57.9%	51.0%	52.2%	25.7%

U.S. Census Bureau (2007b)

The population of the four ceasus tracts that include the study area is most similar to New Mexico overall when comparing income data to the various political entities. As shown in Table 9, the percentage of persons living below the poverty level in 1999 within the four census tracts encompassing the study sees (17.6). poverty level in 1999 within the four census tracts accompaning the study area (1.7.6 percent) was about 1.5 times higher than similar populations in Sants Fe County (12.0 percent) or the City of Sants Fe (12.3 percent) but about the same level as the statewide below-poverty level population.

The per capita income of residents of the study area census tracts in 1999 was only 62 percent of the per capita income of the City of Santa Pe residents and just 67 percent of that of Santa Pe

Table 9. Comparison of selected income data for the State of New Mexico, Santa Fe County, City of Santa Fe, and combined data for Santa Fe County Consus Tracts 8, 12.01, 12.02, and 12.03, which include the study area. Data are based on 1999 income statistics.

			10.00	
Per capita Income*	\$17,261	\$29,594	\$25,464	\$15,651
Persons below poverty level*	18.4%	12.0%	12.8%	17.6%

U.S. Consus Bureau (2007c)
U.S. Consus Bureau (2007d)

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

Figure 41. Land use and ownership features, Camino Alire to Camino Carios Rael.





Figure 42. Land use and ownership features, Camino Carlos Rael to below Siler Road.



Page 48

Figure 43. Land use and ownership features, below Siler Road to below CR 62.




Santa Fe River Study

Page 49


U.S. Army Corps of Engineers, Albuquerque Dist

Figure 44. Land use and ownership features, below CR 62 to Cottonwood Drive.



U.S. Army Corpe of Engineers, Albuquerque District

Figure 45. Land use and ownership features, Cottonwood Drive to NM 599.



Santa Fa River Study

Page 50

Santa Fe River Study

Page 51

The Santa Fe River Trail Corridor Project, a collaborative project between the City and County, is currently underway to develop a continuous trail within the river corridor from Camino Alire to NM 599. According to a City press release (28 March 2007), the river trail under the term of them switch the continuous professor with the continuous continu would be one of three major urban trails in the would be one of three major tuban was as Santa Fe metropolitan area, providing open space, an alternative transportation route, and recreation opportunities for residents and visitors by using the trail to connect a series of parks.

The County does not currently have special zoning or a massagement plan for the river. Their goal is to purchase land or essements from the id or essements from the goal is to purchase iand or essements from too county boundary with the city at Camino Carlos Rael to NM 599 for protection and restoration of the river as well as for recreation uses (C. Pater, Project Manager, Sants Fe County Open Space and Tritle Division, 28 Sept. 2007, pers. comm.). Some projects have already been completed or are courtently in the works, including construction of the new San Ysidro River Park, plans for several new pedestrian and traffic bridges, and a new trail construction project near NM 599.


Currently, recreationists primarily use the study area for walking and related activities (e.g. dog walking, bird watching), jogging, and horseback riding along user-created trails or in the river bed. riding along user-created trails or in the river bed. Walking and jogging use is heaviest in the portion of the study area that is easily accessed on foot from higher-density city-neighborhoods at the east and of the study area. Horseback riding occurs primarily at the western end of the study area where the river bed is more sandy with fewer cobbles and can be accessed from horse properties on the north side of the river.

There is vegetation buffer on either side of the study area for its entire length. In some places, the distance to the nearest building, wall, or other development is as little as 25 feet. In other places, development is as little as 25 feet. In ourse passes, development within a quarter-mile or more is limited to two-track roads, utility lines, and barbed-wire finese. Vegetation is densest and tallest at the eastern end of the study area where the river is restricted to a narrow, entrenched consider. As the river bed widens downstream, vegetation is sparser and lower (i.e. mostly grasses, shrubs, and small trees).

Aesthetic qualities of the study area have been impacted by land-use practices over many decades. These land use practices have included sand and gravel mining, construction of road crossings, dam and diversion construction, sand and gravel mining, construction of road crossings, dam and diversion construction, dumping of solid waste and debris, bank armoring, grading of slopes, and vegetation removal. In more rocest years, off-road vehicle use has detrimentally affected river banks and adjacent land forms by destroying vegetation, breaking down the river banks, and causing crosion and sedimentation. It was not one group of people, a certain time period, or a specific action that caused those changes. It is the cumulative effect of humen activities over several hundred vers., but mostly in the 20° cestary, that hundred years, but mostly in the 20th century, that have brought the Santa Pe River to its current

Along with changes to the natural environment, the biggest aesthetic impacts in the study area are the ubiquitous solid waste debris deposits along the river. Piles of various waste materials have been accumulating over many years,. Some of these solid waste debris accumulations now cover these solid waste deem accumulations down cover several notes along the river banks and adjacent land and extend into the river bed (Figure 46). A fair amount of solid waste debria is still dumped directly into the river but is somewhat less directly into the river but is somewhat less noticeable when flows have moved it downstream. Within the study area are half-buried car bodies and car parts, household appliances, construction

Santa Fe River Study



Homes, property boundary walls, streets, or other structures along this reach are generally set back from the river banks between 25 and 100 feet. from the river banks between 25 and 100 tees. Figure 41 shows that there is generally a buffer strip of vegetation between city streets and the river. A few private parcels with little structural development can still be seen in Figure 41.

At beginning of the study area near Camino Alire, a few mature cottonwoods and class provide shade in the relatively narrow drainage for wildlife, humans, and pets. In some places, tree canopies haman, and pets. In some places, true canopies nearly span the river, providing a sense of privacy and security for river users. Trees and shrubs obscure the walls and fences that border the backyards of residences adjacent to the river and obstact views into and from those yards, creating an additional physical barrier between homewars and recreationists. When the river is flowing, the gurgling water combined with sounds of winds in the trees and birds chiping provides a pleasing respite from the highly-developed urban setting autrounding it. When visited on a warman summer weekday, rounds from nearby homes and streets were surprisingly muffled.

Just a few hundred feet downstream from the Camino Alire crossing, mature trees give way to Just a few hundred feet downstream trom use Camino Alier crossing, nature trees give why to shusts and grasses and the river floodplain wides to that residential properties and recreationists as openly visible, giving the corridor a less "private" feel. Private lands vary from tidy, attractive properties to the less well-kept with dilapidated buildings and old vehicles chattering the sites.

Solid waste debris dumping is evident throughout the length of the study area, particularly at vehicle access locations that are at least partially hidden from public view, such as in arroyes and along dirt roads. Dumping is the most obvious seathers impact in the study area. An example in this segment of the study area. An example in this segment of the study area is a location with piles

of concrete, asphalt, and other construction materials and sumy rusting cer bodies that covers about 700 feet of bank and extends into the river channel (Figure 41).

During visits to the study area in May and June of During visits to the samy sizes in soly and ratio viology, the segment appeared to have the highest amount of recreational use. People using the area were most often sees walking (alone or with a dog) or jogging along the user-created trails that are found on see or both sides of the river this area or in the river bed itself when the flows had recreated.

2.10.2 Camino Carlos Raci to CR 62 (Santa Fe County)

The river segment between Camino Carlos Rael and CR 62 is about 2.6-miles long and covers about 35 percent of the study area. Most of the land in the study area along this segment is either privately owned or owned by Santa Fe County (Figures 42 and 43). Private lands adjacent to the study area are on the north side of the river are commonly used for agricultural purposes, such as horse partures and truck farms.

Agus Fria Street roughly parallels the river to the south, although it is separated from the river by Nouth, although it is separated from the river by private lots (Figures 42 and 43). An Agus Fria is an arterial street in Senia Fe, it is nearly fully-developed commercial and residential interpreted along its length. Commercial enterprises whose properties are adjacent to the south side of the river range from agricultural to light industrial. Some examples of these businesses are a door manufacturing plant, dentist office, plant aussey, runck retails exvice, suate repair, fiquor sales, well-drilling service, landscaping and exceavating services, toofing service, portable toilet supplier, retail window sales, screen printing, accounting service, and a feed supplier. U.S. Army Corps of Engineers, Albuquerque District

waste (e.g. concrete, asphalt, lumber, silt fence, and plastic sheeting), 55-gallon drums, and smaller items normally found in household trush-cans and bottles, paper, plastic groccry bag, and the black trush bags which once held this trush.

The County anti-dumping ordinance has only limited law enforcement backing. The County is trying to limit access to the river as one method combating dumping, and volunteers are often utilized to remove trash from the river (C. Baker. Project Manager, Santa Fe County Open Space and Trails Division, 28 Sept. 2007, pers. comm.)

2.10.1 Camino Alire to Camino Carlos Rael (City of Santa Pe)

The study area within the City's jurisdiction is the approximately 1.5-mile segment from Camino Alire to Camino Carlos Rael and includes about 20 percent of the total river length in the study area (Figure 41). Land ownership along the study area as abown on the City of Santa Fe GIS area as shown on the City of Sai database is primarily as either private area" with a few parcels of city land.

Land uses adjacent to the river in this stretch are primarily residential dwellings and utban streets. There are three city parks along the river: Rio Grande ("pocket") Park, John F. Chiego Park on Pasco do la Conquistadora, and Preschy's Field Park on Agus Fris Street (Figure 41).



Figure 46. Waste piles spilling into the river channel. View is downstream in the rea below Camino Carlos Ruel, 22 June 2007. Photo by J. Pittenger.

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

Since 2000, Santa Fe County has purchased four percels of land (Figures 42 and 43) along this segment of the study area (C. Baker, Project Manager, Santa Fe County Done Space and Trails Division, 28 Sept. 2007, pers. comm.). These properties range between eight and 15 acres in size and activated to both sides of the river for a total length of about 1.1 miles. The sestem-most of these parcels extends both upstream and downstream from the end of Siler Road (Figure 42). County plans for this percel are a new triggen pedestrian bridge to facilitate elimination of the low-water crossing at Camino Carots Rael.

A fifth percel of County land, approximately 70 acres, was formerly federal land snanaged by the U.S. Bureau of Land Management (BLM). This percel was petented in 1975 for "Agas Fris Park" (F. Martinez, Real Estate Specialist, BLM-Taos Field Office, i 10 ctuber 2007, pers. courn.). The land extends upstream and downstream from the CR 62 crossing (Figure 43). Above CR 62, this parcel is on the north side of the river only for 500 foot.

Portions or all of three of these five County parcels are located between the Snn Ysidro crossing and CR 62 (Figures 42 and 43) and form the basis of the newly-created San Ysidro River Park. The County began construction of the park, which stretches for show three-querters of a mile along both sides of the river, in 2005 (C. Baker, 28 Sept. 2007, pers. comm.). Although seartly finished, some rack work near the San Ysidro crossing will be completed in the fall of 2007.

The San Yaidro River Park provides recreational opportunities such as walking, horseback riding, and picaricking. Other recreations use along this study area segments mostly immited to walking or horseback riding in river bed or along occasional

two-track dirt made above the river.

Decades of trush dumping is evident stong the river, particularly on some several undaveloped stretches. One trush dump extends for about 1,000 feet along the north back, covering about three acres. Just one-tenth mile downstream on the same side of the river is sancher dump of encompassing almost two acres (Figure 42; Rocent dumping by businesses along Agua Pris Street whose properties back to the river can be witnessed around the Biler Rods area. It appears that this dumping over the steep south bank (about 20 feet above fits river bed) is covering on Secta Fo County property (Figure 42). Fe County property (Figure 42)

2.1B.3 CR 62 to Cottonwood Drive (Santa Fe County)

The segment from CR 62 to Cottenwood Drive is about 1.5 miles and constitutes about 20 percent of the study area length. Landowsership along segment is primarily private (Figure 44). A small amount of the government-managed lands occur along this segment mear the CR 62 crossing. Below the stressing, County tand purchased from the SLM ancompasses both sides of the river for roughly 0.2 miles. More BLM lands is adjacent to the downstream end of the County land. This purvel is on the north side of the river and extends about 0.2 miles downstream (Naure 44).

Uses of private land on the south side of the river are primarily industrial or commercial with a few homes scattered emong them. Businesses include a septic tank construction company, pawing and appliances salvage years. Sents Fe Baptist Church is a large building on property that is adjacent to the river near South Meadows Road.

The north side of the river is largely undeveloped with the major exception of Cottonwood Mobile Home Park which occupies about 80 acres between the river and NM 99 (Figure 44). The undeveloped land is not in a natural state, however. A 40-acre parcel cast of the mobile home park appears to be used for off road vahiele (ORV) trails, and a fire-scere parcel is completed devoid of vegetation (Figure 44). Many other user-created directions for the strails parallel whis side of the river. Some of these trails parallel this side of the river. Some of these traits parallel the north river bank and are used for access to the river bed, as evidenced in Figure 44.

Santa Fe County has plans for another traffic bridge in this area by extending South Meadows Road from its current terminus on the south side of the river to the north (C.Baker, Project Manager, Santa Fe County Open Space and Trails Divisios, 28 Sept 2007, pers. comm.). This road extension would presumably be to connect to the NM 599 at the CR 62 intersection.

Trush dumps are frequent along the stratch. On the south side of the river, across from the BLM land, a dumping area covers approximately 1.7 acres of private hand adjacent to the river. Send more slightly smaller dumps are on the same side of the river. Many commercial properties appear to use the back area of their lots (i.e. nearest the river) for piling trush as well.

Assilutically, this segment appears "beaten down" by decades of human use. There is little vegetation for acreeiing so dispidated buildings and junked vehicles are easily observed. The partiel with ORV exils is a particular assault on the sances with little to ne vegetation and numerous dirt rails extending over 40 acres, as well as being the source of much noise.

2.10.4 Cottonwood Drive to NM 599 (Santa Fe County)

The western-most segment of the study area includes apportunately! 5-miles of the river. Land in and adjacent to the study area is lawly undeveloped. About 1.2 miles of the river in this segment flows through state land managed by the New Mexico State Land Office (Figure 45). The remaining the state of ining land is private.

Mineral leases records indicate that sand and gravel mining was permitted along this segment by the State Land Office as early as 1928. Currendly, there are no mineral leases along the river in the study area, but there are no number of active commercial leases. Along with several right-of-way leases for roads and utilities in the study area, but State Land Office issued a right-of-way lease in December 2002 for 30 acres to Santh Re Counts for graveries are not seen a lease the Mineral leases records indicate that sand and Fe County for protecting open space along the Santa Fe River. The lease struddles the river bed for about one mile (Figure 45) and is valid for as long as the land is used for the leased purpose.

Two other communical leases for businesses are located in this segment. Both businesses are located on the south side of the river, east of Constellation Drive (Figure 45). The leases are:

- I. 1.5 acres for a bot mix plant and related
- facilities; and

 3.4 acres lease for a construction and trucking

The latter permit, which also includes a residence and related utilities, was issued in 2005 and will expire in 2010. The hot mix plant site has been leased for the same purpose at least since 1987 and has been received regularly since then. The latest lease received was in Cothor 2004 and will expire after five years. These tesses are not very


Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District



grading plugged the restored meander constructed in 2000 on tate Trust land at the downstream end of the study area. Arrows indicate berm that v m that was usked up to block off the sander from the river. View is downstream. Current river channel bank is to the right of the berm, the restored meander channel, now cut-off from the river, is to the left of Pittenger, 24 May 2007.

Figure 47. Upsutherized



Pigure 48. View upstre of the restored meander now isolated from the river. The meander channel is indicated by the channel is indicated by the dotted line. Arrow "A" shows the location of the channel plug shown in Figure 47. Arrow "H" shows the meander cut-off channel created by unauthorized heavy-equipment grading at the site. View is upstream. Photo by J. Pittenger, 24 May 2007.

compatible with plans to develop a recreational trail through the area because of the obronic noise from vehicles and machinery and odors (e.g. disself.tuna, apphall processing) associated with the industrial uses.

Santa Pe County will some productshe out Santa Fo County will soon and retake construction of a new pedestriae bridge and trail along the sand that they have leased from the State Land Office (Figure 45). The bridge will be constructed across the river approximately in line with the end of Countellation Drive. The trail will be built on the north side of the river for about one mile. The County hopes to construct a second pedestriam bridge at the eastern end of this trail and eventually connect it to the Santa Fe River Trail.

In addition, there is a goal of extending the eastern m addition, there is a goal of a standing the eastern territions of the trail north twends NM 599 where it would cross under the highway through existing box culvets and tie into a trail leading to the Caje del Rio Recreation Complex (C. Baker, Project Manager, Santhe Pe County Open Space and Trails Division, 28 Sept. 2007, pers. comm.).

Dumping appears be less provalent along this segment, possibly because much dribris was removed as part of the river restoration project undertaken in 2000 as described in section 1.2. However, anauthorized beavy-equipment grading in the river channel was conducted sometime effer 2000 and the meander created at the restoration. site was intentionally cut off by pushing up a channel plug and a pilot channel was exceavated through the newly-constructed point bar (Figures 47 and 48). The result of this unauthorized grading was abandoument of the meander and loss of an interesting visual feature that created a sense of a natural river channel in the area.

Sente Fe River Study

Page 57

U.S. Army Corps of Engineers, Albuquerque District

3.0 FUTURE CONDITIONS WITHOUT RESTORATION

3.1 Climate

Assuming current trends in human-influenced global climate change continue unchecked, effects in New Mexico will likely include substantially in recoverance will make y include substituting warmer average air temperatures (i.e. increases of 6° to 12° F), higher average minimum temperatures, more frequent episodes of extreme best, fewer episodes of extreme cold, and a longer frost-free period. Effects of global climate chaon precipitation patterns in New Mexico are uncertain and could be either increased or decreased annual precipitation amounts. However, it is likely that snowfall amounts would decrease, spring mow-mest mmost would occur curtim in the year, and more of the total precipitation would occur as rainfall (New

3.2 Groundwater and Surface Water

3.2.1 Hydrology and Water Ouality

Without any restoration projects, stream flow in the Santa Fe River through the study area would continue to occur only sporadically, in response to upstream reservoir releases and storm water runoff. Storm-water nunoff peak flows in the river may increase as more of the watershed is urbanized, particularly in the far northwestern portion of the estchment basin that is undergoing development. Also, climate change natherns may development. Also, climate change patterns may result in increased summer precipitation and less

snow-melt runoff (cf. section 3.1). Under this sconario, sustained spring flow through the study area, similar to that which occurred in 2007, would become more infrequent. On the other hand, short-duration, intense storm-water runoff flows would become more frequent. These conditions would flictly result in less storage in the svaliable shallow althyrium, more bed scour, and less infiltentia into the Themes Exempt and less infiltentia into the Themes Exempt. the available shallow allavium, more bed soor, and less inditation into the Tosque Formation aquifor from the river channel. Surface water quality, during the periods when the river is flowing, will continue to be affected by pollutanas in storm-water runoff. The underlying allavium would continue to be dry throughout most of the study area except for brief periods when it is saturated by infrequent surface water flows in the

3.2.2 Santa Fe River Channel Morphology

The channel of the Santa Fe River in the study area would continue to be influenced by peak flows associated with storm-water ransoff and the condition of grade control structures. Assuming that existing grade control trunchers. Assuming that existing grade control truncher remain in place, the channel of the Santa Fe River through the study channel of the Sants Pe River through the study area should continue to tend toward equilibrium. Some areas, such as the reach below CR 62 (SPR-10) and the reach at and above Prenchy's Park (SPR-2; Plate 6), would continue to widen and evode the steep-walled busks. In other locations, P channel form may continue to adjust to sedament load and discharge to develop C-type channel (meandering) form. However, gives the past record of river management actions it is plausible that additional impacts to the river may occur, which could jolt the system back into a disequilibrium state (e.g. constriction of floodplain or channel width through placement of fills, loss of grade control, by in routing of storm-water flows). Under changes in routing of statin-water interpola-tis scenario, the channel evolution process would be re-set to an early stage and the process of channel adjustment would begin anew.

Grade control structures in the study area are variable in terms of their apparent integrity. The two largest structures, at San Yaidro crossing (GCS-4) and CR 62 (GCS-2; Figure 21) have (OCS-4) and CR 62 (OCS-2; Figure 21) have small seour pools on the downstream side but do not appear to be in danger of becoming undermined in the near future. The San Antonio de Vereda crossing (OCS-5; Plate 6), on the other hand, has a relatively deep scour pool on the downstream side and is crumbling (Figure 22). The long-term persistence of this structure is questionable. The San Antonio de Vereda crossing structure is important in maintaining equilibrium conditions in the upstream reach, which is one of the two main uites in the sudy area where native woody riparian vegetation is becoming established.

Some of the grade coatrol structures downstream from Frenchy's Field appear to be susceptible to undermining and failure. Most notable of these is structure GCS-7 downstream from Camino Carlos Rael (Figure 49). Assuming a bed tipo of about one percent and given the drop below the structure of about three feet, the effect of loss of this structure could potentially extend upstream about 300 feet. The effect would likely persist for many years as the stream channel adjusts its slope and width to the new conditions.

3.3 Air Quality and Noise

If no restoration activities are undertaken in the Santa Fe River study corridor, air quality would not be expected to change measurably. Existing soine levels - human and natural - would also continue at about the same level. The most likely increase in noise levels would come from sounds generated outside of the study area, such as increase traffic on N.M. Highway 599 and city

3.4 Ecological Setting

Development of native riparism regetation in the study area would continue to be limited to those channel areas in or near equilibrium state, where study area would continue to be limited to those channel areas in or near equilibrium state, where deposition of sediment on lateral or point bars occurs and there is sufficient floodplain area to allow the spreading of water and dissipation of energy of flood flows. If existing trends continue, these areas are likely to be limited to the C-type channels segments near Silest Road, the San Ysidro River Park between the San Ysidro rousning and CR 62, and the lower end of the study area from Cottonwood Drive downstream to MM 599. For example, occurrence of sendlings study area from Continuous and the state of seedlings and saplings in the lower reach of the study area, from the NM 599 bridge upstream to Cottonwood Drive indicates that conditions are at least from me NM.399 introduced the conditions are at least periodically suitable for regeneration of cottonwood and willow (Figure 50). Similarly, cottonwood seedlings and suplings and coyote willow seedlings and root-sprouts are common in

Page 60

Seale Fe River Study

U.S. Army Corps of Engineers, Albuquerque Dietrica

Development of native woody riperian vegetation in other segments will continue to be hampered by lack of soil moisture, according peak flows, and expension of non-native invasive species such as expansion of non-native invasive species were skinerian ein, houry-locust, time-of-leaven, and salteedar. Lack of soil moisture, in this context, results from sporadic surface flows combined with fast runoff, very marrow of absent floodplains, and scant deposits of permeable alluvium to absorb

Development of aquatic habitat in the study area is very unlikely in the absence of restoration actions because of the hydrologic and geomorphic trends discussed above. Similarly, wildlife habitat trends discassed above. Similarly, withatte saturation in the study area will contains to be very limited in terms of quality (i.e. the expacity to support a diverse assemblage of native species) and retail extent. Patches of antive woody riparian vegetation will continue to provide the highest quality habitat and these will likely be restricted primarily to the Siter Road vicinity and the lower

Development of suitable habitat for threatened or Development of suitable habitat for threatened or endangared species such as Southwestern Willow Plycatcher in the abrence of restoration is not likely to occur. This is because surface water flow through the study area will continue to be irregular and geomorphic constraints will restrict locations where native woody riparina vegetation will develop. Consequently, evolution of wetlands and dense riparina vegetation in close proximity to persistent surface water (i.e. habitat suitable for Southwestern Willow Flycatcher) is not likely to nocut.

3.5 Socioeconomic Environment

Much of the private property adjacent to the study corridor is already developed for residential, commercial, or industrial uses. Without river restoration activities it is unlikely, therefore, that future socioeconomic conditions in the study r would change substantially

3.6 Land Use

If no ecological restoration action is undertaken in the study area, current trends for land use, recreation, and aestheries would continue. I and uses in the study area would continue to be similar to current uses. It is likely that lands leased by State Land Office for commercial uses (i.e. hot mix plant and construction company) would continue.

These uses would remain as an asselhetic intrusion to the County's new recreation trail in that area. Trash dumps, off-read vehicle impacts to the stream banks and river bod, reeding river banks devoid of vegetation, and a thy river bed with an abundance of visible trash would continue to be asselhed inspairments in the study area, particularly from the perspective of recreationists expecting to enjoy a walk or horreback ride through a natural river environment.

Page 62




Figure 49. Grade control structure GCS-7, downstream from Camino Cartos Rasl, showing failure of the erosion control suren and undermining of the structure. View is upstream. Photo by J. Pittenger, 26 June 2007.




Figure 50. Coyote willow establishment on a point but at the lower end of the study area. View is study area. View II
upstream is a secondary
channel on a point bar that
was restored in 2000.
Photo by J. Pittenger, 20
September 2007.

U.S. Army Corps of Engineers, Albuquerque District

4.0 POTENTIAL NATURAL RESOURCES RESTORATION PROJECTS

4.1 What is Ecological Restoration?

This chapter presents an overview of ecological restoration issues in the study area and potential projects that would contribute to restoring ecological integrity to the river. In the context of this report, restoration is defined as "the process of assisting the recovery of an ecosystem that has been degraded, damaged, or destroyed" (Society of Ecological Restoration, 2004). In this sense, the potential restoration, projects identified below are not intended to maximize neatheric qualities or improve conditions for specific human activities. are not intended to maximize aesthetic qualities or improve conditions for specific human activities. For example, the discussion does not address developing sports facilities or off-road vehicle use areas. It focuses on the ecological integrity of the river condition.

4.2 Restoration Goal and **Objectives**

The need for ecological restoration in the study area is obvious. The river ecosystem has been damaged by flow regulation, channel modification, and establishment of non-antive species. Once the need for restoration is establishmed, it is critical to define a goal and objectives to focus efforts. This is particularly important in an urban/deeni-urban setting such as Santa Fe, where there is strong public interest in restoring the triver. Clear definition of a goal and supporting objectives will go a long way towards resouring unpriver. Clear membron of a good and supporting objectives will go a long way towards coordinating efforts of all stakeholders and prevent parties working at cross-purposes. Establishing a clear goal and objectives also contributes to ensuring success of the restoration program by concentrating efforts where they are

In this light, the following goal and objectives for coological restoration in the study area are offered as a starting point for discussion. The goal is stated as a desired future condition, which provides a narrative picture of where we want restoration to go.

4.2.1 Preliminary Restoration

a desired future condition of the Santa Fe River The desired fishure condition of the Santa Fe River in the study area is a rivorine corridor with natural structure and function, including a stream channel in equilibrium with discharge and estimate supply, flowing surface water duting at least part of the year, and riparism and wetland communities dominated by native plant species.

4.2.2 Preliminary Restoration **Objectives**

Some potential objectives to achieve this goal include the following. The first object, addressing stream flow, is not addressed any further in this report as it is an oegoing effort by the City administration and other interested parties. Similarly, objective number 4, increasing Goodplain area, is a band acquisition issue and is

Manage water supplies to enhance streaflow in the Santa Fe River.

- 2. Reduce flood peaks from storm-water run-off and enhance infiltration of run-off into alluvium.
- Assist natural evolution of the stream channel
- Assist canarate evolution of the stream channel to an equilibrium state.

 Increase floodplain area within the corridor.

 Promote development of riperias and wetland plant communities

4.3 Potential Restoration Strategies and Projects

Using the goals and objectives described above as a guide for actions needed to restore the Sants Fe River in the study area, potential strategies and projects were identified. These strategies and projects were identified. These strategies and projects are grouped in categories that address each of the non-flow related objectives (numbers 2 through 5 above in section 4.2.2). The primary factors limiting ecological restoration in the study area are hydrologic and geomorphic. Consequently, these issues should be addressed first.

4.3.1 Storm-Water Runoff

This restoration strategy, along with the following strategy (4.3.2 Channel Equilibrium) are the two primary restoration actions needed in the study area. The aim of the storm-water runoff strategy is to slow down run-off, reduce flood peaks, and increase indiffration of storm water into the shallow altuvium of the Santa Pe River.

Potential restoration projects in the study area that are consistent with this strategy deal with detention of storm-water flows at outfall points within the corridor (cf. Plate 6). However, it should also be recognized that perhaps a more important set of restoration projects should be implemented in the larger watershed area. Examples of the latter types of projects include: capturing run-off from roof areas on individual properties and directing the water into Prench drains or similar structures; replacing impervious paving with porous aurfacing; construction seepage basing along arroys; and installing check dams in head-cutting guillies (Riley, 1998: 340-354).

The major arroyo confluence points or storm-water outfalls in the study area (cf. Plate 6) that lend themselves to creation of retention basins

- 1. SW-1, the Arroyo Toureon confluence (Station 366+00);

- SW-1, the Arroyo Torreon confluence (Station 366-00);
 SW-2, the storm-water drainage confluence at Frenchy's Park (302-61);
 SW-9, the arroyo confluence near Nix Stables; (Station 290-72);
 SW-1, Storm drain outfall below the Siler Road dead end (Station 253-62);
 SW-1, storm drain outfall near the intersection of South Meadows Road and Ams Eris Streng (Station 99-88);
- Agua Fria Street (Station 99-88);

 6. SW-19, stown-water channel confluence (Station 56+00); and

 7. SW-20, NM 599 stown-water drainage
- channel confluence (Station 12+45).

Retention basins would serve as excellent sites for establishing stands of coyote willow and cottonwood or improving existing stands by retaining run-off and allowing it to percolate into the alluvium (Figures 51 and 52). Development of wettands dominated by bettacous plants (e.g. acque, bullaubat, rusbee, obligate and facultative wetland forbs) may also be possible if retention basins are not subject to excessive sediment descriptions.

Santa Fe River Study

Page 64



Important design considerations include expected run-off volumes, water velocity, sediment loading, and grade courted. These persuncters should be carefully assessed and incorporated into design. A conceptual design for storm-water retention A conceptual design for sto basins is shown in Figure 53.

The potential for creation of a wetland with herbacous plants would be greatest at the Arroyo de la Joya conflaines as Frenchy's Park (SW-2, Plate 6). An existing excerned pond area could be reconstructed at this site by enlarging it significantly, grading the excerned area to configure soveral sub-basins, and directing stormwater flows from the deniange into the wetland and then from the wetland into the river.

An impervious liner was installed in the existing dayreasion; this should be removed to allow retained storm-water to infiltrate into the allowium. Willow, cottonwood, and firebrocous plants should be planted in the securety area. Anticipated storm-water mooff volume, sediment input, and scour characteristics should be evaluated carefully and incorporated into the evaluated carefully and incorporated into the estign. If sediment londing is expected to be substantial, one or more settling sub-basins should be planted. These would likely require periodic insistenance to remove accumulated material and maintain sediment-rapping capacity. The resulting storm-water detention basin may have standing water for sustained periods, similar to the cuisting storm-water detention pond just upstream from the NM 599 bridge on the left terrace (Figure 54).




Figure 53. Conceptual design of a storm-water




Figure 51. Storm drain outfall SW-12 at Siler Road. The culvert outfall (A) is shadowed by (A) is anadowed by vegetation. Cottonwood asplings (B) and coyote willow (C) are growing amongst Siberian elm and rebbitbrush at the site. Photo by K. Yori, 12 July



Figure 52. Stand of structure SW-5. View is natream. Photo by J. enger, 15 June 2007.

Santa Fe River Study






Figure 54. Stottn-water detention pond ne the NM 599 bridge. The pond held water (photo A, 24 May 2007) for a considerable period in spring 2007. Also water invols receded, the bottom of the pond was colonic reason in spring 2007. After water levels recorded, the bottom of the pond was colonis by smussi plasets (8, 19 September 2007). Woody vagetation on the pond banks inclusionyste willow and Rio Grande cottonwood. Photoh by J. Pittunger.

4.3.2 Channel Equilibrium

Establishing channel equilibrium conditions in the study area is a vital prerequisite for effective coulsgical restoration. As long as channel of degradation and channel widening are dominant geomorphic processes, woody riparian vegetation cannot become established at persist at a site.

Potential restoration projects that apply to this strategy address developing an understanding of current conditions and rectifying obvious factors

contributing to disequifibrium in the study area. The former action is of utmost importance. The services of a probasional geomorphologist should be employed in a comprehensive assessment of the classmel in the study truck, including a controlled survey of the longitudinal profile and an appropriate number of cross sections. In particular, evaluation of classmel evaluation stage, current condition of grade controls, need for modification or replacement of controls, and softment manaport characteristics should be thoroughly addressed.

Potential channel restoration projects that could be implemented now include reconnecting the restored meander at the downstream end of the study area and installation of a novel grade control structure in the Frenchy's Park reach.

As discussed in section 2.10.4, a mea As discussed in section 2.10 A, a meanaber restored in 2000 between stations 39-10 and 49-00 was cut-off by unsusthorized grading and excavation in the river channel sometime after 2000 (Figures 47 and 48). The shandcased meanaber could be reconnected to the river by removing the berms at the indet and sutlet of the channel, filling the cut-off channel, and restoring the large point bar that had been created at the site by placement of fill, predign antiver hards and control and co

Another project that may be considered is construction of a new grade control structure to replace existing structures GCS-10 and GCS-11 at Frenchy's Park (between stations 302+00 and 306+00, Plate 6).

This could be implemented in conjunction with construction of a storm-water detention basin at the site (cf. section 4.3.2). The conceptual design the site (cf. section 4.3.2). The conceptual design is to replace the existing concrete and gabion grade control structures with sish structures that emulate in appearance and function the outcropping of relatively impermeable Tesaque Formation red conglomeratic silts (cf. Figure 8). The intent would be to restore a shallow alluvial er table and conditions similar to what may water table and conditions similar weathing have been present at historic Cioneguits Spring (see discussion on pages 9-10). The intent would be for the structure to serve as a grade control and be for the structure to serve as a grame control sail also to retain shallow groundwater moving through the alluvium, which would be charge at the site by the storm-water input from the detention basin (Figure 55). The crest of the structure sill should be placed at a sufficient elevation to cause deposition of sediment scurcure still should be placed at a sufficient elevation to cause deposition of sediment (aggradation) upstream because the bed has accured down to relatively impermeable strata and storage volume for shallow groundwater is limited.




Figure 55. Conceptual design for ted conglomerate outcrop grade simulated conglomerate outcrop grade-control structure. The structure would also serve to retain shallow groundwater moving through the coarse alluvium, potentially bringing it to the surface during wet periods.

Santa Fa River Study

Page 68

U.S. Army Corps of Engineers, Albuquerque District

5.0 PREPARATION, COORDINATION, AND CONSULTATION

5.1 Preparation

John Pittenger, Senior Ecologist Elue Earth Ecological Consultants, Inc.

Karen Yori, Senior Planner Blue Barth Ecological Consultants, Inc.

5.2 Coordination and Consultation

College Baker, Project Manager Santa Fe County Open Space and Trails Division

Gilbert Burregn, Youth Works Consultant City of Santa Pe

Mary Rilen Carroll, Minerals Management Analyst State Land Office, Oil, Gus and Minerals Division

David Chapman, (Acting River Coordinator)

Francina Martinez, Realty Specialist Bureau of Land Management, Taos Field Office

U.S. Army Corps of Engineers, Albuquerque District

Major channel restoration work should not be conducted in the study area until a thorough geomorphic analysis is conducted. This is protections are to the reach downstream from CR 62, which is deeply incised and unstable. Channel restoration should be carefully planted using regional curves and a suitable reference site using resjocal curves and a suitable reference site to determine average values and range for inhibity in design criteria for morphological features such as channel slope, basefull width and depth, meander wavelength, belt width. For example, the restoration project at the downstream end of the study area involved developing a regional curves for the relationship between drainage area and channel morphology persuants gaged and un-paged sites in the Sania Fe County area (Blue Earth Boological Consultant, Lec., unpubl. dats). A reference reach was Inc., unpubl. data). A reference reach was identified and field measurements were made to develop design criteria. These included a channel alope of 0.73 percent, average bankfull width of 1.55, fiest, average bankfull depth of 1.27 fiest, and sincosity of 1.54, in a project reach that had sincosity of 1.54, in a project reach that had channel bed material with a D_o of 1.2 mm (Blue Barth Boological Consultants, Inc., uspubl. data). Use of inappropriate materials in (a.g. harp boulders) and methods (a.g. construction of overwidened channel) should be avoided. Inc., unpubl. data). A reference reach was identified and field measurements were made to

4.3.3 Native Riparian Vegetation

This strategy is directed at increasing the extent of This strategy is directed at increasing one accent on native riparain vegetation in the study area. Two main actions can be undertaken that would contribute to this objective. Parts, supplemental planting of native woody riparian species can be implemented in areas that are in equilibrium condition. Planting in unstable reaches should be for the planting in unstable reaches should be consistent with geomorphic factors influencing stability area addressed (e.g. Briggs, 1996: 100-101). Second, removal of non-native species can be conducted and the plants replaced with native

species. The areas of focus for supplemental planting actions should be 1) the Siler Road area and 2) the downstream end of the study area, where restoration work conducted in 2000 resulted in dramatic increases in cover to contonwood sad willow. The Siler Road area is a good candidate for the initial focus, as it appears to have good potential as indicated by natural recruitment of cottonwood and willow.

Planting of coltonwood and coyote willow can be most easily accomplished by using dormant pole or whip eatings (e.g. Briggs, 1996: 71-77). Planting techniques should follow guidance developed by the Naharal Resources Conservation Service, Plant Materials Center in Los Lunas developed by the Nahmal Resources Conservation Service, Plant Materials Center in Los Lunas (Appendix B). A rotary harmore filted with a ose-inch dismeter, three-foot long bit or a backhoe fitted with a "stinger" on the excavator arm are suitable methods for sugaring deep hotes for plusting whips or small-dismeter poles. These methods should work well in the soil types found

4.3.4 Community Involvement

As an end note, the need to involve the community in restoration planning and implementation cannot be stressed snough. Through involvement in removing trust from the river, planting vegetation, removing one-native plants, monitoring river conditions, and other activities, community sevaruous of the value of a living river increases. Through involvement with restoration, the computative claims as more claims as living river increases. Through involvement win-restoration, the commenting pains as appreciation of place and the river becomes a meaningful part of their lives. Community interest in restoration can be fostered by communicating beastin such as enhancing neighborhoods, preserving history and culture, increasing recreational opportunities, creating jobs, and providing unique educational opportunities (Riley, 1998: 12-22).

U.S. Army Corps of Engineers, Albuquerque District

6.0 REFERENCES

- Allred, K. 2006. A Working Index of New Maxico Vascular Plant Names. http://web.amsu.odu/-kallred/ hattweb/Working%20lndms-tile.htm.
- nerican Sivers. 2007. America's Most Endangered Rivers, 2007. http://www.americanzivers.org/site/ PagoSorver/pagoname=AR7_MBR2007
- Briggs, M. K., 1996. Riparian Scoaystem Recovery in Arid Lands: Strategies and References. The University of Arizonu Press, Tucson, Arizona.
- City of Santa Fe. 2007. Santa Fe Trende 2007. City of Santa Fe Planning and Land Use Department, Banta
- Dunne, T. and L. B. Leopold. 1978. Water in Environmental Planning. W.H. Prommen and Company, New
- Francessa, N. M. and D. W. Johnson. 1946. Physiographic Divisions of the Conterminous U.S., scale 1:7,660,000. U.S. Geological Survey, Washington, D.C.
- FOTURRA, R. T. T., D. Sperting, J. A. Biscorete, A. P. Clevenger, C. D. Cutaball, V. H. Dale, L. Fehrig, R. France, C. R. Goldman, K. Hennate, J. A. Jones, F. J. Swenton, T. Turrentine, and T. C. Winter. 2003. Road Ecology, Science and Solutions. Island Press, Washington, Covolo, London.
- Grant, P. 2002. Santa Fe River Watershed Restoration Action Strategy. Santa Fe Watership Santa Fe, New Mexico.
- Hárvey, M. D. and C. C. Watson. 1986. Fluvial processes and morphological thresholds in incised channel restoration. Water Resources Bulletin 22(3): 359-168.
- Heggen, R. J. 1997. Leavons Learned. Channel Stabilization in Santa Fe, New Mexico. Unpublished report prepared for the U.S. Army Corps of Engineers, Albuquerque District, Albuquerque, New Mexico.
- Hink, V. D. and R. D. Ohmset. 1984. Middle Ria Grande Biological Survey, Final Report. Report prepared for the U.S. Army Corps of Baginsers, Albuquerque District, in fulfillment of contract no. DACW47-81-C-0015.
- Konig, D. J., S. D. Connell, F. J. Pazzagtia, and W. C. McIntoth. 2002. Radefinition of the Ancha Formation and Phacana. Plaistocoma denocition in the Santa Fe ambayment, north-central New Mexico. New Mexico. and Phiocene-Plaistoce Geology 24(3): 75-87.

- Kottek, M., J. Grisser, C. Beck, B. Rudolf, and F. Rubel. 2006. World Map of the Köppen-Geiger climate classification updated. Meteorologiache Zettschrift 15: 259-263.
- ttiewski, F. E. 1995. Geologic synopsis of La Villa Real de la Santa Fe de San Francisco de Assisi. Pages 2-4 (n: Bauer, P. W., B. S. Kuns, N. W. Dunbar, K. E. Karlstrom, and B. Harrison (eds.). Geology of the Santa Fe Region, New Mexico. New Mexico Geological Society Guidebook, 46th Field Conference, September 17-10, 1995
- Lange, C. 1998. Clean Water Act Section 404 Permit Application for the Santa Fe River Channel Trail and Greenway Improvements Project, Santa Fe County, New Mexico. City of Santa Fe Public Works Department, Engineering Project Division, Santa Fe, New Mexico. 4 February 1998.
- maras, J. and P. Drakos. 1995. Geohydrologic characteristics and hydrocarbon contamination of the shallow alluvial/Tenque Formation aquifer, Santa Fe, New Mexico. Pages 307-311 in: Bauer, P. W., B. S. Kues, N. W. Dunbur, K. E. Karlstron, and B. Harrison (eds.). Geology of the Santa Fe Region. New Mexico Geological Society, Forty-Sixth Annual Field Conference, September 27-30, 1995.
- Leopeld, L. B. 1994. A View of the River. Harvard University Press, Cambridge, Massach
- Leopold, L. B. and J. P. Miller. 1956. Ephemeral Streams Hydraulic Factors and Their Relation to the Drainage Net. U.S. Geological Survey, Geological Survey Professional Paper 282-A.
- Leopald, L. B., M. O. Wolman, and J. P. Miller. 1992. Fluvial Processes in Geomorphology. Dover
- Lewis, A. C. and F. West. 1995. Conceptual hydrologic systems for Santa Fe County. Pages 299-306 in: Bauer, P. W. B. S. Kues, N. W. Dunbar, K. E. Karlstrom, and B. Harrison (eds.). Geology of the Santa Fe Region. New Mexico Geological Society, Porty-Sixth Annual Field Conference, September 27-30, 1905.
- Makepeace, D. K., D. W. Smith, and S. J. Stanley. 1995. Urban stormwater quality: summicontaminant data. Critical Reviews in Environmental Science and Technology 25(2): 93-139.
- MeAda, D. P. and M. Wasiolek. 1988. Simulation of the Regional Geohydrology of the Texaque Aquifer System Near Santa Fe, New Mexico. U.S. Goological Survey Water-Resources Investigations Report 87-
- ional Climatic Data Center. 2007. Climate Monitoring Reports and Products. U.S. Department of Commerce, National Oceanic and Atmospheric Administration Satellite and Information Service. http://www.ncdc.nosa.gov/en/climate/research/monitoring.html

U.S. Army Corps of Engineers, Albuquerque District

- U.S. Census Burens, 2007a. American Fact Finder. P7. Ruce (8) universe: total pagai Census 2000 summary file 1(8F 1) 100-percent data...http://factfinder/census.gov/
- U.S. Cemens Bureau, 2007b. American Fact Finder. PS. Hupanic or Latino by race [17] universe: total population; data set: Cemens 2000 summary file 1(SF 1) 100-percent data. http://factfinder/census.gov/
- U.S. Census Bureau. 2007c. American Fact Finder. P82. Per capita income in 1999 (dollar) [1] universe: total population; data set: Census 2000 summary file 3 (SF 3) sample data. http://factfinder.census.gov/
- U.S. Ceases Baress. 2007d. American Fact Finder. P87. Poverty status in 1999 by age [17] universe population for whom poverty status is determined; data set: Census 2000 summary file 3 (SF 3) sample data. http://factfinder.census.gov/
- U.S. Census Bureau. 2007e. American Fact Finder. T1 Population estimates [9]; data set: 2006 population estimates. http://factfinder.census.gov/
- U.S. Environmental Protection Agency. 2007. Air data. Non-attainment Areas Map Criteria Air Poliutants. http://www.spa.gov/six/dats/reports.html
- Western Regional Climate Center. 2007. New Mexico Climate Summaries for Cooperator Stations.

 Western Regional Climate Center, National Oceanic and Atmospheric Administration, Reno, Nevada.

 http://www.wrcc.dri.odu/indox.html
- W.J. Miller Engineers, Inc. 2000. Santa Fe River Demonstration Project, State Trust Reach, Sediment Transpartation and Hydraulics Analysis. Prepared by William J. Miller Engineers, Inc. and Blue Earth Ecological Consultants, Inc. for the New Mexico State Land Office, 1 May 2000.

U.S. Army Corps of Engineers, Albuquerque District

- Natural Resources Commervation Service. Web Soil Survey. Soil Survey Staff, Natural Res Conservation Service, U.S. Department of Agriculture. http://wobsoilsaurvey.arcs.usda.gov/
- New Mexico Chimate Center. 2007. Climate of New Maxico. Chimate Contex, New Maxico State University,
 Las Cruces, New Moxico. http://weather.nameu.odu/
- New Maxine Environment Department. 2005. Potential Effects of Climate Change on New Idaxico.

 Agency Technical Work Group, State of New Mexico. http://www.nmenv.state.mm.us/aqh/cc/Potential_

 Effects_Climate_Change_NM.pdf
- w Mexico Environment Department. 2007. 2006-2008 State of New Mexico Integrated Clean Water Act \$103(6)/\$305(6) Report. New Mexico Environment Department, Surface Water Quality Bureau, Santa Pe, New Mexico.
- Oragon Climate Service. 2007. Comparative Climate, New Mexico. Oragon fints University, Corvellis, Oragon. http://www.ocs.orst.edu/page_links/comparative_climate/new_maxica/new_mexico/new_mexico/new_
- Pitt, R., R. Field, M. Lalor, and M. Brown. 1995. Urban stormwat and treatability. Water Environment Research 67(3):260-275.
- Rend, A. S., D. J. Konig, G. A. Smith, S. Raiser, J. Rogers, and P. W. Bauer. 2000. Preliminary Geologic Map of the Santa Fe 7.5-minute Quadrungla. New Mexico Bureau of Geology, New Mexico Tech, Socorro, New Mexico.
- Riley, A. L. 1998. Restoring Streams in Cities, A Guide for Plannara, Policymakers, and Citizens. Island Press, Washington, D.C. and Covela, California.
- Rosern, D. L. 1994, A classification of natural rivers. Catena 22(3): 169-199.
- Rosgen, D. L. 1996. Applied River Morphology. Wildland Hydrology, Pagosa Springs, Colorado.
- Shrehn, R. R., R. A. Thompson, S. A. Mizor, V. J. S. Gsuch, and T. R. Brundt. 2005. Geologic Map of the Agua Fria Quandrangie, Santa Fe County, New Mexico. U.S. Geological Survey Scientific Investigations Map 2896.
- Society for Ecological Restoration. 2004. The SER International Primer on Ecological Restoration for Boological Restorational, Science and Policy Working Group, Tucson, Arken
- Splegel, Z. and B. Baldwin. 1963. Geology and Water Resources of the Santa Fe Area, New Mexico. U.S. Geological Survey Water-Supply Paper 1525.

Santa Fe River Study

U.S. Army Corps of Engineers, Albuquerque District

APPENDIX A

Animal Species Potentially Occurring in Riparian or Aquatic Habitats in Santa Fe County

The following list of vertebrate animal species potentially occurring in riperian and aquatic habitats was generated from a questy of the Biota information System of New Messice (BISOMM) database, which is maintained by the New Mexico Department of Game and Faits and is located on the world-wide work is wow, bisser-m.org. The database was accessed on 14 November 2007 and was queried using the following

County Name = Santa Fo Habitat = Aquatic 9 ---County rease—Sease re Habitat — Aquatic; Semi-Aquatic; Fully Aquatic; Riperisa Gap Veg — Lowland Rhurian (cottouwood/sycamore); Aquatic: Riverino/Lacu

CONTROL NAME	SERVICE LAVE	
AMPHIBIANS (4 species)		
tiger salamender	Ambyatoma tigrinum	
Great Plains toad	Bufo cognetus	
Woodhouse's toed	Bufo woodhouell	
свлуоп tree frag	Hyla amelicolor	
REPTILES (23 species)		
onsale box turtle	Terrapene ornate	
collered lizard	Crotephytue collerie	
leopard Hzard	Gamballa wiałzonii	
losser dariesa tzard	Holbrookle maculete	
roundtail horned lizard	Phrynozoma modestum	
prairie lizard	Sceloporus undulatus	
northam tree Hzard	Uroseurus ornetus	
Chihuahuan spotted whiptell	Aspidocella exsengula	
Colorade checkered whiptuil	Aspidoscells tesselete	
Plateau spotted whilpts!	Aspidocelle velox	
many-lined skink	Eumecee multivirgatus epipleurotis	
Great Plains skink	Eumeces obsoletus	
glossy snake	Arizona elegane	
com snake	Elaphe guillets	
westem hognose snake	Heterodon nesicus	
night anaka	Hypsiglene torquete	
desert kingsnake	Lampropeltis getula spiendide	
milk enake	Lampropeitis trianguium celaenops	
amooth green snake	Opheodrys vernalis blanchardi	
coachwhip	Masticophia Sagolium	
desert striped whipenake	Masticophia teenistus teenistus	

Senta Fe River Study Page 76

U.S. Army Corps of Engineers, Albuquerque District

Yellow-billed Cuskeo	Coccygus americanus occidentalis
Greater Roadrunner	Geococcyx californianus
Barn Owl	Tyto albe pretincola
Long-eared Owl	Asio plus
Flammufeled Owl	Otuz Sammeckus
Western Screech Owl	Otuz kennicotti
Great-hamed Owl	Subo virginianus
Mexican Spotted Owl	Strix accidentells lucide
Northern Pygmy Owl	Glaucidium gnoma californicum
Burrowing Owl	Athene cunicularie hypugues
Common Nighthawk	Chordelles minor
Belted Kingfisher	Ceryle alcyon
Black-chinned Hummingbird	Archillochus alexandri
Broad-bried Hummingbird	Seisephorus platyoerous platycerous
Rufous Hummingbird	Selasphorus rufus
Northern Flicker	Coleptee suratus
Lewis's Woodpecker	Melanurpsa lewis
Downy Woodpacker	Picoldes pubescens leucurus
Helry Woodpecker	Plaeides villosus
Ladder-backed Woodpacker	Picoldes scalaris
Willemson's Sepsucker	Sphyrapicus thyroideus netalies
Red-naped Sapeucker	Sphyrapicus nuchalis
Olive-sided Flycatcher	Contopus acaperi
Western Wood-Pierre	Contopus sordidulus
Southwestern Willow Flycetcher	Empidonex trelf8 extimus
Dusky Flycatcher	Empidonax oberholauri
Cordilleran Flycalcher	Empidonax occidentalis

Processor Wiles, 18	
gopher scake	Pluophis centler
wastem bisckneck garter enake	Themnophia cyrtopeus cyrtopeis
wandering garter anake	Themnophia elegens
checkered garter anake	Themophia mardenus marclatus
New Mexico garter enake	Thammophie sirtalis dorenila
western diarnondback rattlesnake	Crotake strox
BIRDS (143 species)	
Double-created Cormorant	Phalacrocorex euritus
Great Blue Heron	Ardea harodies herodies
Wood Duck	Alu apones
Turkey Vulture	Cathertes aure
Northern Harrier	Circus cysneus hudeonius
Cooper's Hewk	Apolpitar cooperii
Sharp-shinned Hawk	Accipitur atriatus velox
Northern Gostawk	Accipiter gentitis
Zone-tailed Hawk	Buteo albonoletus
Red-telled Hereft	Suteo jamelcentie
Sweineur's Hawk	Buteo sweinsoni
Ferruginous Hawk	Buteo regalis
Baid Eagle	Heliseatus leucocaphalus alastanus
American Kestrel	Falco sparverius sparverius
Peregrine Fatcon	Falco peragrinus anatum
Sandfelli Crane	Grus canadensis
Spotted Sendpiper	Actilis menuturia
Rock Dove	Columba livis
Band-tailed Pigeon	Columbe festiets fasciate
Mourning Dove	Zenelde mecroure

Sania Fo River Study Page 7

U.S. Army Corps of Engineers, Albuquerque District

R. D. Lee and C. Lee a	The second se
Black Phoebe	Sayomis nigricens semistre
Say's Phoebs	Seyumle saya
Ash-throated Plycatcher	Mylerchus cinerascene cincerascene
Casalria Kingbird	Tyrannus vociforans vociforans
Western Kingblid	Tyransare varificalle
Loggerhand Shrike	(,anius iudovicianus
Northern Shrike	Lanius excubitor invictus
Casaln's Vireo	Vireo cassinii
Plumbeovs Virso	Vireo plumineus
Solitary Vireo	Virea eofterius
Warbling Vireo	Virse glion amintonii
Blue Jey	Cymnocitta oriztata bromia
Steller's Jay	Cyanopite stated metrolophe
Western Scrub Jay	Aphelocome californics
Black-billed Magple	Pice hudeonie
American Crow	Corvus brackyrhynckos
Common Ravan	Corvus corex sinustre
Tree Swellow	Techyolosta biolor
Violet-grape Swallow	Tachycineta thelessina (apide
Northern Rough-winged Swallow	Steigidupteryx semipernis semipennis
Barn Swallow	Hrunda rustice srythrogester
Citi Swallow	Petrochelidan pyrrhanota
Juniper Titmouse	Secolophus ridgwsyl
Black-capped Chickedee	Poecile atricepille
Mountain Chickadee	Poecile gambeli gambeli
Buehdt	Psakriparus minimus
White-browning Nethatch	Sitta carolinanale nelsoni

Santa Fe River Study Page 78 Santa Fe River Study Page

COMMON NAME	a example and
Pygmy Nuthalch	Sitte pygmaea melanotia
Red-breasted Nuthatch	Sitte cenedensis
Brown Creeper	Corthia emericana
Sewick's Wran	i hryomanez Dewiciol
House Wren	Troglodytes aedon parkmarnii
Canyon Wren	Catherpea mexicanus conspensus
Golden-crowned Kinglet	Regulus satrapa
Ruby-crowned Kinglet	Regulus colondule colondule
Blue-gray Gnatcatcher	Polioptila caerulea emoeniesima
Mountain Bluebird	Statia ourrecoldes
Western Bluebird	Slalle mexicana bairdi
Townsend's Solkaire	Mysdeales townsendii (omnsendii
American Robin	Turdus migratorius
Swainson's Thrush	Catherus ustulatus
Hermit Thrush	Catherus guitatus
Gray Cathird	Dumetalla carolinansis rulloriusa
Northern Mockingbird	Mimus polygiotics leucopterus
Sage Thrusher	Oreoscoptes montanus
American Pipit	Anthus rubescens
Cedar Waxwing	Sombycille cedrorum
European Sturling	Sturrus vulgeris
Orange-crowned Warbler	Vermivore celete
Virginia's Warbler	Vernivora virginiae
Nashdie Warbler	Vermirora ruficapilla ridgereyi
Yellow Warbler	Dendroice patechia
Townsend's Warbler	Dendroice townsendi
Black-throated Gray Warbler	Dendroica nigrescens

Santa Fe River Study Page 80

Golden-crowned Sparrow	Zonotrichie asricapille
White-throated Sparrow	Zonotrichia albicolla
Derk-eyed Junco	Junco hyemele
Bullock's Orlole	fcterus bultockii
Baltimore Oniole	fotorus palbule
Scatt's Origin	foturus perfaorum
Red-winged Blackbird	Ageleius phoeniceus
Brewer's Blackbird	Esphagus cyanoceptatus
Common Grackle	Quisicalus quiscula versicolor
Great-tailed Grackie	Quiniculus mesicanus
Brown-headed Cowbird	Mointhrus ager
Evening Grosbeak	Coccothraustes respertinus
House Finch	Carpodacus mexicanus frontalis
Lesser Goldfinch	Cardunia peatria paatria
American Goldfinch	Cardunila triatis pallitius
Pine Siskin	Certuels pinus pinus
MAMMALS (57 apocios)	
Virginia oposeem	Didelphie virginiene virginiene
dusky shrew	Sorex monticulus
southwestern myotis	Myofs auriculus apachs
fringed myotie	Myolia thysanodes thysanodes
long-legged myotis	Alyatis valens interior
western small-footed myotis	Myotis cilialebrum melenarisinus
big brown bat	Eptesious fuscus palidus
honry but	Lesiurus cinereus cinereus
pale Townsend's big-eared bat	Corynorkinus townsendii pallescena
palled bet	Antrozona palikus palikus

La Comballe Aug.	
Black-litrosted Green Warbler	Dendroica virena virana
Yellow-rumped Warbler	Dendroice coronate
Grace's Warbler	Dendroice gracies gracies
Bay-breasted Warbler	Dendroka cestanea
Macgillivray's Warbler	Operantis tolonial
Common Yallowthroat	Geothlypie kiches
Wilson's Warbler	Wilsonia puellis
Yellow-breasted Chat	Ictoria virans auricollia
Western Tanager	Pirange Iudoviciene
Hepatic Tanager	Piranga flava
Indigo Bunting	Passerina cyanes
Lazvi Bunting	Passerina emoene
Blue Grosbeak	Phouticus caerules interfuss
Black-headed Grosbeak	Phoutique melanocaphatus
Rose-breasted Grosbenk	Pheuticus Indovicienus
Canyon Townse	Pipilo Necus
Spotted Towne	Plpilo meculatus
Green-tailed Townse	Plpilo chiorurus
Chipping Sparrow	Spizete personne erizonee
Brewer's Sparrow	Spizella brewerl
Clay-colored Sparrow	Spizula pallida
Lark Sparrow	Chondestes grammecus strigatus
Vesper Sparrow	Pagecates graminous
Song Sparrow	Meloepiza melodia
Lincoln's Sparrow	Melospiza (Incolnii
Harris's Sparrow	Zonotrichia querulu
White-crowned Sparrow	Zanotrichia laucophrys

Sents Fe River Study Page 81

U.S. Army Corps of Engineers, Albuquerque District

desert cottontall rabbit	Sylvilagus audubonii
black-tailed jack rabbit	Lepus celliomicus
spotted ground equirmi	Spermophilus spilosoms
rock equirmi	Spermophilus variegatus grammunus
Bolta's pocket gopher	Thomosoye bettee
silky packet mouse	Perognethus llevus
Ord's kungaroo rat	Dipodomys ordii
white-throsted woodrat	Neotoma albigule
baqvar	Cestor caredensis
western harvest issues	Relithrodontomys megalistile
deer mouse	Peromyaqua meniculatus
white-footed mouse	Реготувана (висорие
brush mouse	Peromystus bay@ rowley!
northern grasshopper mause	Onyohomys leucogaeter
house mouse	Mus mustulus
common porcupine	Erethizon derestum
coyole	Canile febrane
gray lox	Urocyon cimeroergenteus scottii
black bear	Uraus americanus ambiyospa
ringtali	Betseriscus eskulus
comitton recoon	Protyen latar
American badger	Textidue tuxus
western apotted akunk	Spilogale gracilie
striped skuris	Maphitis maphitis
mesmbin lien	Fells conscior
bobcat	Lynx rufus ballayi
mude deer	Odocolleus bernionus

Senta Fe River Study Page 82 State Fe River Study Page 83

Deep Planting

APPENDIX B

Guidelines for Riparian Planting


New Mexico Plant Materials Center Brochure #7105, Quidelines for Planting Dommet Whip Cuttings to Revegetate and Stabilize Stream Banks, available on-line at: http://www.plant-materials.arcs.usda.gow/pubs/ampmcbr7105.pdf

New Mexico Plant Materials Center Brochure #7106, Quidelines for Planting Longstem Transplants for Riparian Restoration in the Southwest, available on-line at: http://www.plant-materials.urcs.uada.gow/pubs/nmpmcbr7106.pdf

Senta Fe River Study

Page 84





Page 88

MCDONALD, MELISSA A.

From:

John Buchser <jbuchser@comcast.net>

Sent:

Friday, September 02, 2016 10:06 AM

To:

MCDONALD, MELISSA A.

Cc:

Andy Otto; Phil Bove

Subject:

River Commission - Priority Setting and annual community meeting

Fellow River commissioners,

For our October meeting (10/13) I would like to spend a significant part of the meeting discussing our priorities for the next 2-3 years.

Please provide a prioritized list of about 10 to 15 items to Melissa no later than Monday prior to the October meeting, then we can merge ideas together and discuss as a group. If you can provide a rough idea of cost and possible funding source (if funding is needed) that would be very helpful. If there are recommendations from subcommittees, please submit those as well.

It has been several years since we have had a community meeting. I propose that we plan on a community meeting, roughly in the May 2017 timeframe, potentially replacing our regular monthly meeting. I'd like to start a discussion about this after our priority setting. This would be a good opportunity for us to share what our vision and priorities for the River with the Mayor, Councilors, and the press.

Thanks,

John

