Herbicide	Semera	Do not irrigate one hour before or after application;Do not apply more than 24 ounces per acre per year; Do not apply to saturated soils or prior to a rain event.	annual/perennial weeds pre and post emergent	<u>View Label</u>	<u>View MSDS</u>	59639-120-91234	flumioxazin 51%	non crop areas turf and ornamental	Caution
Herbicide	Zaltys	Do not apply more than 24 ounces per acre per year; Do not reapply within 30 days.	annual/perennial weeds pre and post emergent	<u>View Label</u>	<u>View MSDS</u>	59639- 120-91234	flumioxazin 51%	right of way, medians parking and storage areas, fence line,airports bare ground non- crop areas	
Herbicide	Civitas WeedFREE	Do not irrigate 24 hours after application Do not apply prior to rain Spray when weeds are young and actively growing Delay mowing 1-2 days after application	Broad leaf weeds	<u>View Label</u>	<u>View MSDS</u>	69526-15	2,4-D dimethylamine salt 0.885% Mecoprop-p, dimethylamine salt 0.573% Dicamba, dimethylamine salt 0.109%	Putting Greens and other improved turf sites.	Caution
Fungicide	Subdue MAXX	Do not apply more than the maximun season total for the active ingredient as stated on the label of the product containing the lowest seasonal total on that crop.	damping off and root-stem disease and foliar disease organisms	<u>View Label</u>	<u>View MSDS</u>	100-796	Mefenoxam 22%	Ornamental, greenhouse lanscape, non- fruit bearing contaner grown plants	Caution
Fungicide	Actinovate SP	Apply prior to disease onset; follow practices known to reduce disease development	Soil borne and foliar diseases	<u>View Label</u>	<u>View MSDS</u>	73314-1	Streptomyces lydicus WYEC 108 0.0371%	Greenhouse, nursery landscape, interirscape ornamentals,transplant dip soil drench	Caution
Insecticide	AzoSol	Medium Rate For preventative to medium infestations when pests are present.) 2-3 tsp/1000 sq. ft.	Control of whiteflies, scale insect and caterpillar larvae, bark beetles.	<u>View Label</u>	<u>View MSDS</u>	81899-4	Azidiracthin 6%	Nursery, greenhouse, landscape plants, trees	Caution
Fungicide	Phospho-Jet	For terrestrial use only	For control of Root rot, Phytopthora, Anthracnose disease organisms	<u>View Label</u>	<u>View MSDS</u>	74578-3	Phosphoric acid 45.8%	Landscape (Deciduous and Pine) trees.	Caution
Herbicide	Rodeo	No Soil Activity: Weeds must be emerged at the time of application to be controlled by this product.	For control of herbaceous and woody vegetation as foliar or cut stump treatment.	<u>View Label</u>	<u>View MSDS</u>	62719-324	Glyphosate isopropylamine salt 53.8%	Rights of way, industrial sites, railroads, airports, recreational areas, rangeland, other non-crop sites	Caution
Herbicide	Garlon 4 Ultra	Do not apply directly to water, to areas where surface water is present	Control of woody plants and annual and perennial broadleaf weeds in non-crop areas	<u>View Label</u>	<u>View MSDS</u>	62719-527	triclopyr 60.45%	Cut stump application on non-crop areas including rights of way, industrial sites, and roadways.	Caution
		This list is not inclusive of all pesticides that may	be required to handle specific emergency weed and pest o	occurrences as pe	er City of Santa Fe	Integrated Pest Management	(IPM) Ordinance. 2001-10-7.10.E		